REVIEW

Role of adipokines in embryo implantation

Davoud Jafari-Gharabaghloo^{1,2}, Mostafa Vaghari-Tabari², Hajar Oghbaei³, Laura Lotz⁴, Reza Zarezadeh², Yeganeh Rastgar Rezaei⁵, Mahnaz Ranjkesh⁶, Mohammad Nouri^{2,7}, Amir Fattahi^{4,8}, Saba Nikanfar^{2,8} and Ralf Dittrich⁴

¹Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

²Department of Clinical Biochemistry and Laboratory Medicine, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

³Department of Physiology, Tabriz University of Medical Sciences, Tabriz, Iran

⁴Department of Obstetrics and Gynecology, Erlangen University Hospital, Friedrich-Alexander University of Erlangen–Nürnberg, Erlangen, Germany

⁵Department of Medical Biotechnology, Faculty of Advanced Medical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran

⁶Medical Radiation Science Research Group (MRSRG), Tabriz University of Medical Sciences, Tabriz, Iran

⁷Department of Reproductive Biology, Faculty of Advanced Medical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran

⁸Stem Cell Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Correspondence should be addressed to A Fattahi or S Nikanfar: amirfattahi@gmail.com or nikansaba@gmail.com

Abstract

Embryo implantation is a complex process in which multiple molecules acting together under strict regulation. Studies showed the production of various adipokines and their receptors in the embryo and uterus, where they can influence the maternal-fetal transmission of metabolites and embryo implantation. Therefore, these cytokines have opened a novel area of study in the field of embryo–maternal crosstalk during early pregnancy. In this respect, the involvement of adipokines has been widely reported in the regulation of both physiological and pathological aspects of the implantation process. However, the information about the role of some recently identified adipokines is limited. This review aims to highlight the role of various adipokines in embryo–maternal interactions, endometrial receptivity, and embryo implantation, as well as the underlying molecular mechanisms.

Key Words

- ▶ adipokine
- ▶ embryo implantation
- ▶ endometrium
- ▶ pregnancy
- ▶ trophoblast

Endocrine Connections
(2021) **10**, R267–R278

Introduction

Nowadays, many advances have been achieved in the field of assisted reproductive technology (ART) and this technique could help many infertile couples to have their children. However, the main problem with this technology is the low rate of implantation after the embryo transfer (1). Statistics showed that more than 50% of embryos could not be implanted after transferring into the uterus and even in some patients the implantation failure can be seen after several transfers of good quality embryos which are described as repeated implantation failure (2).

Implantation is one of the most important steps toward pregnancy initiation, through which blastocyst invades the epithelium of the endometrium (3). To successful implantation, both embryo and endometrium should be met in a precise time and place, which is recognized as the ‘implantation window’ (3).

During this window, the endometrium is completely ready for receiving the blastocyst. In a normal menstrual cycle, this period is from days 16 to 22; this would be approximately 5–10 days after the luteinizing hormone (LH) surge (4). The embryo implantation contains three steps, at the first stage, the blastocyst attaches with a loose connection to the implantation site in the uterine (opposition), and then trophoblasts attach to the epithelium of the endometrium (adhesion), and finally, at the last stage, the cells invade to the stromal site of the endometrium (invasion). All the steps should be precisely regulated to make the successful implantation possible (5).

Various factors are involved in the implantation process such as cytokines (e.g. IL1, IL6, and leukemia inhibitory factor), prostaglandins (e.g. PGE2), growth factors (e.g. epidermal growth factor (EGF), vascular endothelial

growth factor (VEGF), and heparin-binding EGF), matrix-degrading enzymes (e.g. matrix metalloproteinases (MMPs)), matrix-degrading enzymes inhibitors (e.g. tissue inhibitor of metalloproteinase (TIMPs)) and molecules that have a role in cellular adhesion (e.g. integrins, selectins, and cadherins) (2, 3). It has been documented that adipose tissue-derived factors can also affect implantation (6, 7). Adipose tissue acts as an endocrine tissue and secretes hormone-like factors called adipokines. Adipokines are a kind of cytokine that contributes to energy metabolism, inflammation, immunity, and angiogenesis, as well as, reproductive maturity and fertility (6, 7, 8, 9). Regarding the latter effect, it has been well documented that obesity or excess fat can negatively influence female fertility and ovarian function such as what we see in obese women with polycystic ovary syndrome (PCOS) (10, 11). The role of adipokines has also been shown in embryo implantation (6, 7). Owing to the importance of adipokines in fertility and implantation and the lack of comprehensive study on the involvement of adipokines in embryo implantation, in this review, we aimed to address the roles of adipokines in embryo implantation and the underlying mechanisms.

Adipokines in embryo implantation

Many studies have shown the presence of adipokines and their receptors in the woman's reproductive systems emphasizing their role in female fertility. Adipokines are also present in the uterus and placenta where they can influence implantation and pregnancy as well as maternal-fetal transmission of metabolites (6, 7). It has been demonstrated that various stages of implantation are profoundly regulated by steroids, including progesterone, estradiol, and androgens. On the other hand, the role of adipokines in the regulation of the secretion of gonadotropin-releasing hormone (GnRH), gonadotropins, and steroids suggesting their involvement in the implantation process (12). However, there is no consensus regarding the effects of adipokines on hormones which can be due to different reasons such as species differences, the dose of adipokines (13, 14), size and stage of ovarian follicles (15), and phase of the menstrual cycle (16). The association of adipokines with steroidogenesis and hormones has previously been reviewed in depth by our group (17). Interestingly, adipokines can be under the regulation of hormones. For example, the stimulatory effects of estrogen, testosterone, LH, and follicle-stimulating hormone (FSH) on resistin expression have been demonstrated in ovarian follicles (18). Moreover, it has been reported that estradiol

could induce adiponectin expression (19). However, the inhibitory effects of steroids have also been reported on adipokines in ovarian follicles (19).

It has been documented that alteration of adipokine levels may result in different female reproductive issues. For example, both abnormal decreased and increased levels of adipokines have been reported in gestational diabetes mellitus, PCOS, preeclampsia, endometriosis, and intrauterine growth retardation; however, some studies did not find an association between adipokine levels and gynecological issues (12). In addition, it has been demonstrated that increased levels of adipokines can alter their functions and may cause adipokine resistance (20). Adipokines can link energy metabolism and reproduction. Implantation and pregnancy are intensely reliant on the balance of energy (21, 22), and therefore metabolic abnormalities can cause implantation failure (23). The roles of different adipokines in implantation and pregnancy have been mentioned by previous studies. In this regard, various adipokines such as leptin, adiponectin, apelin, chemerin, progranulin, retinol-binding protein 4 (RBP4), and visfatin have been investigated (7, 24, 25, 26, 27). In the following sections, we explain the adipokines and their possible role in embryo implantation. Moreover, the role of adipokines in implantation and possible underlying mechanisms are summarized in Table 1.

Leptin

Leptin is a 16 kDa polypeptide hormone encoded by the obese (*OB*) gene and mostly produced by white adipose tissue. It acts through the leptin receptor that mainly transmits signals through the Janus kinase/signal transducer and activator of transcription signaling pathway. The main role of leptin in the body is regulation of energy balance by affecting cellular metabolism and appetite (28, 29). Interestingly, leptin can act as a pro-inflammatory cytokine due to its structural similarity with IL6. In this regard, leptin increases the expression of inflammatory cytokines such as tumor necrosis factor α (TNF α) and IL6 (30, 31). Leptin is also associated with the adaptive immune system and can enhance the proliferation and survival of T cells (30). Studies have well documented that leptin is involved in the reproductive system. For instance, this adipokine is essential for the onset of puberty, and also can affect the hypothalamus-pituitary-gonadal axis (32).

In the female reproductive system, it has been demonstrated that leptin receptor is present on the surface of granulosa and theca cells, and stimulates the production of the steroids in these cells (33). In supporting the role of

Table 1 Role of adipokines in embryo implantation-related functions and their mechanism of action.

Adipokine	Function	Mechanism of action	Ref
Leptin	Remodeling of the endometrial epithelium (human)	Stimulating proliferation and enhancing Fas ligand-induced apoptosis in EECs	(39)
	Repairing endometrial epithelium after embryo implantation (human)	ND	(39)
	Inducing blastocysts adhesion (mice)	Induction of αv and $\beta 3$ integrin in EECs	(40)
	Stimulating trophoblast invasion (human, mouse)	Upregulating MMP expression in trophoblast cells	(123, 124)
	Induction of implantation-related inflammation (human)	Stimulating the production of IL6 and chemokines in EECs and ESCs	(42)
Adiponectin	Inhibition of decidualization (human)	ND	(41)
	Playing a role in maternal recognition of pregnancy and implantation (porcine)	Upregulation of basal and insulin-stimulated steroidogenic enzymes, including HSD3B, StAR, and CYP11A1 and secretion of A_4 and P_4 by the endometrium and myometrium	(51)
	Involving in the development of the preimplantation embryo and uterine receptivity (mouse)	Activation of autocrine and paracrine mechanisms	(56)
	Involving in decidualization		
	Increasing uterine receptivity	Stimulation of AKT/PI3K and MAPK signaling pathways	(52)
	Inducing proliferation and attenuating apoptosis of uterine luminal epithelial cells (porcine)		
	Involving in implantation and regulating endometrial inflammation (human)	Increasing phosphorylation of AMP-activated protein kinase in ESCs and EECs	(50)
Apelin	Regulating luteal phase of the estrous cycle (pig)	Decreasing IL1 β -induced secretion of IL6, IL8, and MCP1 from ESCs	(64)
	Regulating early placental development (human)	Stimulating P_4 secretion and HSD3B activity in the middle of the luteal phase	(67)
	Decreasing the uterine contractions and facilitating embryo implantation (human)	Enhancing proliferation of trophoblast cells via APJ and ERK1/2, Stat3 and AMPK α signaling	(69, 70)
	Exerting hypotensive effects on blood vessels (rats)	ND	
	Inducing the uterus myometrium contraction (rats)	Stimulating the phosphorylation of eNOS, activation of guanyl cyclase, and cGMP production in endothelial cells	
Chemerin	Maintaining early pregnancy and preventing embryo abortion (human)	Partially via protein kinase C pathway	(81)
	Regulating endometrial receptivity and trophoblast invasiveness (porcine)	Phosphorylation of ERK1/2	(25)
	Involving in maternal recognition of pregnancy (porcine)	ND	(25)
	Decidualization and vascular remodeling during early pregnancy (human)	Possibly via the conceptus-secreted substances, including uPA and E2	(83)
	Promoting uterine receptivity and implantation (human, mice)	Accumulating NK cells at the implantation site	(68, 82)
	Increasing trophoblast invasion (human)	Augmenting AMPK α activity	(83, 84)
Visfatin	Involving in implantation (mice)	Production of IL8 and IP10 by NK cells	(99)
	Affecting uterine immune responses and morphological structure (rat)	Increasing PCNA expression and maintaining the balance between apoptotic and anti-apoptotic elements in the uterus	(97)
	Involving in the regulation of the placental angiogenesis (human)	Modulating inflammatory responses by regulating the expression of eosinophil, myeloperoxidase, and inflammatory cytokines (IL1 β , IL6, and TNF α)	(100)
RBP4	Increasing trophoblast proliferation and invasion (human)	ND	(105)
		Suppressing PI3K/AKT signaling pathway and upregulating MMP2 and MMP9	

(Continued)

Table 1 Continued.

Adipokine	Function	Mechanism of action	Ref
Progranulin	Stimulating blastocyst growth and adhesion (mouse)	ND	(118)
	Involving in decidualization (murine)	ND	(119)
	Establishing placenta by involvement in hypertrophy and proliferation of the endometrial epithelium, growth and migration of cytotrophoblast, and maternal angiogenesis (mink)	ND	(26)
	Playing role in uterine angiogenesis and facilitating the implantation (human)	ND	(121)
	Involving in blastocysts hatching (mouse)	ND	(118)

A4, androstenedione; AMP, adenosine monophosphate; AMPK, AMP-activated protein kinase; APJ, apelin receptor; HSD3B, 3beta-hydroxysteroid dehydrogenase; cGMP, cyclic guanosine monophosphate; CYP11A1, cytochrome P450 11A1; E2, estradiol; EEC, endometrial epithelial cells; ERK1/2, extracellular signal-regulated kinases 1 and 2; eNOS, endothelial nitric oxide synthase; ESC, endometrial stromal cells; MAPK, mitogen-activated protein kinase; MCP-1, monocyte chemoattractant protein 1; MMP, matrix metalloproteinases; ND, not defined; NK, natural killer; P4, progesterone; PCNA, proliferating cell nuclear antigen; PI3K/AKT, phosphatidylinositol-3-kinase/protein kinase B; StAR, steroidogenic acute regulatory protein; STAT3, signal transducer and activator of transcription 3; TNF, tumor necrosis factor; uPA, urokinase plasminogen activator.

leptin in embryo implantation, expression of its receptor (OBR) has been observed on the endometrial cells and it has been reported that decreased expression of the receptor is associated with subfertility (34). Leptin is locally produced by endometrium and blastocysts showing its importance in embryo-endometrium crosstalk during implantation (35). Findings from subsequent studies further reinforced the potential roles of leptin in the pregnancy process. In this regard, it has been shown that injecting recombinant leptin for 8 days resulted in pregnancy in leptin-deficient female mice suffering from infertility. More interestingly, pregnancy did not occur when leptin administration was stopped 0.5 or 3.5 days after mating, but it occurred when the treatment stopped 6.5 or 14.5 days after mating (36). Given that the implantation initiates 5 days after mating in mice, it seems that the most important role of leptin in pregnancy is related to the implantation stage. Another study showed that the expression of leptin receptors in the endometrium was significantly increased during the luteal phase, the phase of receptivity (37). A study also reported that the blocking of leptin receptors on the third day of pregnancy could impair embryo implantation in mice (38). Also, studies have demonstrated that leptin is involved in the proliferation and apoptosis of uterine epithelial cells, uterine receptivity, uterine immune system, and decidualization, which are necessary for implantation (39, 40, 41). On the other hand, the expression of leptin could only be detected at the blastocyst stage during the preimplantation period, the stage that can be implanted (37).

Regarding the underlying mechanisms of leptin action in implantation, it has been documented that leptin could significantly increase the adhesion rate of mice blastocysts

via inducing the expression of adhesion molecules such as beta 3 integrin in endometrial epithelial cells (40) (Fig. 1). Leptin has also an association with the endometrial immune system and can induce the expression of several implantation-related inflammatory cytokines in the endometrium. In this regard, it has been indicated that leptin treatment could increase the expression of IL6, IL8, GRO α , monocyte chemoattractant protein-1, and macrophage inflammatory protein-3 α (MIP3 α) in the endometrial epithelial and stromal cells (42). On the other hand, the essential role of inflammation in the embryo implantation process has been well-documented (43). Leptin is also involved in decidualization, a necessary step for pregnancy initiation (41). However, the effect of leptin on the decidualization is inhibitory, and therefore, extra levels of leptin as seen in the endometriosis patients can also be involved in implantation failure (41).

Adiponectin

Adiponectin is produced by the adipose tissue, liver, bone, and placenta (44). Adiponectin acts through two receptors, AdipoR1 and AdipoR2. AdipoR1 transduces adiponectin signal through AMP-activated kinase (AMPK), whereas AdipoR2 activates peroxisome proliferator-activated receptor alpha (PPAR α) (45). Adiponectin plays an important role in the regulation of energy metabolism via increasing insulin sensitivity, glucose uptake, and fatty acid oxidation, and also attenuating gluconeogenesis (46). Adiponectin is also involved in the immune system and appears to have anti-inflammatory properties via attenuating the nuclear factor-kB signaling pathway and subsequently the production of IL6 and TNF α in

Figure 1

Schematic diagram demonstrating the involvement of adipokines during different stages of the blastocyst implantation in the endometrium. ZP, zona pellucida; BL, blastocyst; CT, cytotrophoblast; ST, syncytiotrophoblast; ES, endometrial stroma; EE, endometrial epithelium.

macrophages. Roles of adiponectin in the reproductive system have also been indicated. Adiponectin attenuates LH secretion from the pituitary gland and regulates GnRH receptor expression (47). A study reported that mice lacking adiponectin showed folliculogenesis issues and impaired fertility potential (48). It has also been demonstrated that the serum levels of adiponectin were directly correlated with the number of retrieved oocytes in women who underwent in vitro fertilization (49).

The expression of adiponectin receptors in the endometrium and the involvement of adiponectin in the embryo implantation-related processes encouraged scientists to investigate the potential roles of this adipokine in embryo implantation. In this respect, it has been found that *ADIPOR1* and *ADIPOR2* were expressed in epithelial and stromal cells of the endometrium and their expression was markedly increased during the mid-luteal phase, the time of embryo implantation (50). Low expression of adiponectin receptors has also been reported in the endometrium of women with unexplained recurrent implantation failure (7). The production of steroids is important for appropriate embryo implantation and it has been shown that adiponectin enhances the expression of steroidogenic enzymes such as 3 β -hydroxysteroid dehydrogenase and steroidogenic acute regulatory protein (51). Furthermore, adiponectin induces proliferation and attenuates apoptosis of uterine luminal epithelial cells, and subsequently increases uterine receptivity. These effects are mediated by the stimulatory effect of

adiponectin on phosphatidylinositol-3-kinase/protein kinase B (AKT/PI3K) and MAPK signaling pathways (52). Adiponectin can also induce endometrium receptivity via reducing IL6 secretion by endometrial epithelial cells (50, 53). Adiponectin can decrease nitric oxide (NO) levels in the endometrium. Given that in the pre- and peri-implantation stages the expression of inducible nitric oxide synthase (iNOS) is increased in the endometrium, it can be postulated that adiponectin plays a regulatory role in NO-related processes during the implantation (54, 55). However, further studies are needed to clarify how adiponectin acts in favor of implantation by reducing NO synthesis, a factor that is required for implantation. Interestingly, the expression of adiponectin receptors is increased during decidualization, suggesting the potential role of this adipokine in decidualization (56). To sum up, adiponectin is involved in embryo implantation via affecting steroidogenesis, the proliferation of uterine epithelial cells, regulating endometrial inflammation, NO synthesis, and endometrium receptivity, and decidualization; however, future studies are likely to reveal further insight into the role of adiponectin in embryo implantation.

Apelin

Apelin is encoded by the *APLN* gene as a preproprotein and following the post-translational modifications, several active forms of apelin are produced, including apelin 36,

apelin 17, and apelin13, among which apelin13 has the highest biological activity. It acts through a G protein receptor named APJ (57, 58, 59). Apelin is involved in many biological functions, including angiogenesis, blood pressure regulation, heart contraction, water intake, and anti-inflammatory processes (57, 60). Apelin also has direct and indirect roles in the reproductive system. In this regard, it has been reported that apelin 13 could attenuate the secretion of LH, FSH, and prolactin from the pituitary gland (61). Apelin and its receptor are expressed in oocytes and follicles and their expression has a positive correlation with ovarian follicle growth (62, 63). These findings support the important role of apelin in female fertility.

Apelin can be involved in the embryo implantation process via enhancing the basal steroid secretion in ovarian cells and inhibiting FSH-induced steroid secretion. Moreover, apelin stimulates progesterone secretion in the middle of the luteal phase and also stimulates hydroxysteroid dehydrogenase (HSD) activity (64). There is emerging evidence that the expression level of apelin increases during the estrous cycle and decreases following corpus luteum degeneration (65). Apelin is also expressed by endometrial tissue and its expression is more pronounced during the secretory phase of the menstrual cycle, confirming the possible role of apelin in endometrial receptivity and implantation (66). In this respect, it has been seen that apelin induces some of the important implantation-related signaling pathways, especially MAPK signaling (67). It seems that the association of apelin with different signaling pathways and their effect on implantation is an interesting issue that deserves further study. For example, apelin has recently been shown to enhance the proliferation of trophoblast cells by phosphorylation of extracellular signal-regulated kinase (ERK1/2), Stat3, and AMPK α (67). On the other hand, it has been shown that the ablation of the *Prkaa1* and *Prkaa2* genes which are involved in encoding the catalytic domains $\alpha 1$ and $\alpha 2$ of AMPK α , could impair embryo implantation in mice (68). Therefore, apelin may play an important role in implantation by affecting AMPK α signaling. Moreover, apelin can weaken uterine contractions and consequently facilitate embryo implantation in the human myometrium (69). In contrast, in a study on rats, it has been reported that apelin induced uterus contraction partially via the protein kinase C pathway (70). Additionally, apelin has antioxidant properties and could increase catalase activity and attenuate reactive oxygen species production (71, 72). Since oxidative stress could have detrimental effects on embryo implantation (73, 74, 75), the apelin-induced antioxidant defenses could support successful implantation.

Given the role of apelin in modulating the immune system and inflammation, and on the other hand, the importance of these events in embryo implantation, it is likely that apelin has further roles in the implantation process. However, few studies have been performed on the association of apelin with implantation and further investigations are required to clarify the underlying mechanisms.

Chemerin

Chemerin is highly expressed in white adipose tissue, liver, and lung. Moreover, its receptor, chemokine-like receptor 1 (CMKLR1), is abundantly found in immune cells (76). This adipokine is involved in the differentiation of adipose cells, chemotaxis of immune cells, angiogenesis, and production of cytokines such as IL6 (76, 77). Chemerin exerts an inhibitory effect on ovarian follicular development and can induce arresting of follicle growth and apoptosis of granulosa cells (78). In addition, studies have shown that chemerin attenuated FSH-induced follicular steroidogenesis which raised the possibility of involvement in PCOS (79).

The endometrium, placenta, trophoblasts, and conceptuses express members of the chemerin system which may reflect the role of this adipokine in embryo-maternal interactions (80). Furthermore, it has been shown that local factors in the uterus could affect the protein expression of the chemerin system during early gestation (25). It has been indicated that this adipokine can prevent embryo abortion via regulating ERK1/2 phosphorylation, an important signaling pathway in embryo implantation (25, 81). In this respect, an upregulation of CMKLR1 in the decidua and a reduction of chemerin in plasma have been reported in women who had spontaneous abortions (81). In a study on pigs, it has been seen that the protein expression of chemerin and CCRL2 (C-C motif chemokine receptor-like 2; a chemerin receptor) in the endometrium was in the highest levels during implantation (25). Moreover, this group demonstrated that the levels of GPR1 (protein-coupled receptor 1c; a chemerin receptor) were significantly increased in the endometrium during embryo migration. An enhanced expression of chemerin and its receptors (CMKLR1, CCRL2, and GPR1) was also observed in the myometrium of pigs during embryo implantation (25). An elevated expression of chemerin has been found in the stromal cells during decidualization (81). Chemerin can augment AMPK α activity which is involved in uterine receptivity and implantation (68, 82). This adipokine also plays a key

role in the accumulation of natural killer (NK) cells at the implantation site (83); the NK cells induce trophoblast invasion by producing different cytokines such as IL8 and IFN γ -inducible protein 10 (84). Moreover, chemerin takes part in vascular remodeling and angiogenesis across early pregnancy (83).

The clinical evidence has shown that elevated proinflammatory factors such as MIP1 β , TNF α , and dendritic cells (DCs) following endometrial biopsy could increase the rate of successful implantation (85). Chemerin is involved in the regulation of inflammatory processes (86). Chemerin is also able to enhance the accumulation of DCs at the site of inflammation (87). Moreover, a correlation has been reported between serum levels of chemerin and TNF α (88). Interestingly, Gnainsky *et al.* (89) found a positive association between levels of TNF α in the endometrium and successful implantation. Nevertheless, chemerin also has anti-inflammatory and pro-oxidating effects (82, 90) that may influence embryo implantation; so further studies are needed to clarify the exact role of chemerin in implantation.

Visfatin

Visfatin or nicotinamide phosphoribosyltransferase (NAMPT) functionally acts as an enzyme restricting the conversion of nicotinamide to nicotinamide mononucleotide (NMN) which subsequently activates the synthesis of nicotinamide adenine dinucleotide (NAD $^{+}$) and thus plays a regulatory role in the energy metabolism (91, 92). Visfatin is also involved in inflammatory responses (secretion of IL1 β , IL6, and TNF α), prevention of neutrophils apoptosis, and maturation of B cells (93, 94). Moreover, recent studies have shown that visfatin could bind and activate the Toll-like receptor 4 (95) and also due to its insulin-mimetic activity, can bind the insulin receptor and cause insulin resistance and consequently regulate glucose levels (96).

Based on the immunohistochemical findings, visfatin is expressed in the myometrium, perimetrium, and especially the endometrium, showing the potential role of this adipokine in functions of the uterus (97). Interestingly, the expression of this adipokine is under the control of steroids as estrogen increases visfatin levels and progesterone has an inhibitory effect on its expression (98). Annie *et al.* (99) have reported that visfatin can be involved in implantation through increasing proliferating cell nuclear antigen expression and maintaining the balance between apoptotic and anti-apoptotic elements in the mice uterus.

As another mechanism, Yang *et al.* (97) have shown that visfatin modulates inflammatory responses by regulating the expression of eosinophil, myeloperoxidase, and inflammatory cytokines and therefore plays a vital role in the uterine immune responses. In line with this study, Kim *et al.* (100) have also demonstrated that visfatin is involved in the regulation of the placental inflammatory response and angiogenesis. Although the information regarding the role and the molecular mechanism of visfatin in embryo implantation is not adequate, the findings of several studies elucidate its involvement in early pregnancy (101).

RBP4

Vitamin A is indispensable for the function of the immune system, epithelial tissue maintenance, and differentiation. Besides, it plays an undeniable role in the reproductive system including ovarian follicle development and endometrial receptivity (102, 103). RBP4, a component of adipokines (104), acts as a specific carrier of vitamin A from the liver to the peripheral tissues and is involved in the regulation of cell differentiation and invasion (105, 106). There are different types of RBP (1 to 4) that have diverse functions in the various tissues; RBP4 is the most abundant type in the serum. Interestingly, in addition to the liver, this protein is produced by the endometrium (107). It has been demonstrated that the plasma concentration of RBP4 is regulated by progesterone and it reaches the maximum level at the mid to late phases of a menstrual cycle to provide vitamin A for the uterus (108, 109). Given the importance of vitamin A in uterine functions, the availability of this vitamin in the uterus is contingent on the concentration of RBP4. The immunocytochemistry analysis has verified the secretion of RBP4 by the uterus (107), which infers the importance of RBP4 in uterine functions. More interestingly, studies have indicated that the expression of RBP4 is increased on days 7 and 13 of a cycle, which are two important checkpoints for preimplantation (110). The HSD17B is involved in the regulation of estrogen and androgen levels and also plays an important role in decidualization. Studies on human endometrial stromal cells have shown that RBP4 could induce 17 β -HSD expression in the epithelial cells (111, 112). Elevated levels of RBP4 during the decidualization can confirm that RBP4 is involved in decidualization via inducing HSD17B expression. Also, RBP4 can induce the secretion of VEGF, an essential factor for angiogenesis and implantation (113). A study by Li *et al.* (105) showed that overexpression of RBP4 could increase trophoblastic

proliferation and invasion via suppressing the PI3K/AKT signaling pathway and upregulating the matrix-degrading enzymes (MMP2 and MMP9), which are important in the implantation and decidualization processes.

Progranulin

Progranulin (PGRN) structurally acts as a precursor of granulin, epithelin, and PC cell-derived growth factor (26, 114). In the intact form, PGRN has anti-inflammatory effects but it acts as a proinflammatory factor when it is cleaved into granulin (115). PGRN is expressed in almost all tissues but it can abundantly be found in rapidly dividing tissues such as keratinocyte, enterocyte, gastrointestinal tract, and uterine epithelium. PGRN is highly expressed in the uterine epithelium during implantation (116). Furthermore, the expression of PGRN has been reported in preimplantation embryos with the highest levels in the blastocyst stage (117). In addition to blastocyst outgrowth, PGRN can promote the adhesion of blastocyst (118). The maternal and embryonic expression of PGRN suggests its role in the implantation and placentation process. Moreover, the involvement of PGRN in endometrial decidualization has been demonstrated (119). In mink, the association of PGRN expression with hypertrophy and proliferation of endometrial epithelium as well as growth and migration of cytotrophoblast suggests its role in embryo implantation and placentation (26). Given the anti-inflammatory properties of PGRN and its role in the growth of epithelial cells and also the presence of this adipokine in all stages before the implantation, it can be postulated that PGRN is involved in embryo implantation. It has been shown that the expression of PGRN dramatically drops following implantation (120). Perez *et al.* (121) showed that PGRN could play an effective role in uterine angiogenesis and facilitating implantation in humans. It has also been documented that the expression of PGRN is increased during human embryo implantation, and dysregulated PGRN expression is associated with pregnancy-related diseases such as abnormal placental angiogenesis (122). Qin *et al.* (118) have indicated that PGRN is secreted from the mammalian blastocysts into the surrounding medium and it is involved in the preparation of the blastocyst for hatching and implantation. In this regard, they reported that adding exogenous PGRN to the culture medium could considerably increase blastocyst hatching, an essential process for implantation (118). Overall, according to the results of studies, PGRN might be an effective factor for

embryo implantation, however, to clarify the exact role of this adipokine in implantation further studies are required.

Conclusion

Adipokines and their receptors are present in embryonic and uterine components suggesting their significant roles in fetal-maternal crosstalk during implantation. The involvement of adipokines has been demonstrated in the regulation of peri-implantation embryo development, trophoblast activity, endometrial receptivity, decidualization, and implantation. Moreover, abnormal alterations in the levels of adipokines may result in pregnancy complications, including implantation failure and recurrent spontaneous abortions (7, 41). These alterations might be either as a consequence of disturbances in the upstream factors, such as hormones or as a causative factor in affecting downstream implantation-related pathways, for example, steroidogenesis. However, the exact underlying mechanisms by which adipokines are involved in implantation are not thoroughly understood and merits further investigation. Furthermore, additional comprehensive studies are required to determine whether the measurement of adipokine system components in uterine fluid during early pregnancy can provide valuable data regarding the prediction of successful implantation.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was supported by the Stem Cell Research Center of Tabriz University of Medical Sciences, Tabriz, Iran (grant number: 62615).

Acknowledgments

This article forms part of the PhD thesis of Saba Nikanfar.

References

- Somigliana E, Vigano P, Busnelli A, Paffoni A, Vegetti W & Vercellini P. Repeated implantation failure at the crossroad between statistics, clinics and over-diagnosis. *Reproductive Biomedicine Online* 2018 **36** 32–38. (<https://doi.org/10.1016/j.rbmo.2017.09.012>)
- Lédée N, Petitbarat M, Chevrier L, Vitoux D, Vezmar K, Rahmati M, Dubanchet S, Gahéry H, Bensussan A & Chaouat G. The uterine immune profile may help women with repeated unexplained embryo implantation failure after *in vitro* fertilization. *American Journal of Reproductive Immunology* 2016 **75** 388–401. (<https://doi.org/10.1111/aji.12483>)

- 3 Namiki T, Ito J & Kashiwazaki N. Molecular mechanisms of embryonic implantation in mammals: lessons from the gene manipulation of mice. *Reproductive Medicine and Biology* 2018 **17** 331–342. (<https://doi.org/10.1002/rmb2.12103>)
- 4 Thiagarajan DK, Basit H & Jeanmonod R. *Physiology, Menstrual Cycle*. Treasure Island, FL, USA: StatPearls, 2019. (available at: <https://www.ncbi.nlm.nih.gov/books/NBK500020/>)
- 5 Ochoa-Bernal MA & Fazleabas AT. Physiologic events of embryo implantation and decidualization in human and non-human primates. *International Journal of Molecular Sciences* 2020 **21** 1973. (<https://doi.org/10.3390/ijms21061973>)
- 6 Reverchon M, Ramé C, Bertoldo M & Dupont J. Adipokines and the female reproductive tract. *International Journal of Endocrinology* 2014 **2014** 232454. (<https://doi.org/10.1155/2014/232454>)
- 7 Dos Santos E, Serazin V, Morvan C, Torre A, Wainer R, de Mazancourt P & Dieudonné MN. Adiponectin and leptin systems in human endometrium during window of implantation. *Fertility and Sterility* 2012 **97** 771.e1–778.e1. (<https://doi.org/10.1016/j.fertnstert.2011.12.042>)
- 8 Lago F, Dieguez C, Gómez-Reino J & Gualillo O. The emerging role of adipokines as mediators of inflammation and immune responses. *Cytokine and Growth Factor Reviews* 2007 **18** 313–325. (<https://doi.org/10.1016/j.cytogfr.2007.04.007>)
- 9 Christodoulatos GS, Spyrou N, Kadillari J, Psallida S & Dalamaga M. The role of adipokines in breast cancer: current evidence and perspectives. *Current Obesity Reports* 2019 **8** 413–433. (<https://doi.org/10.1007/s13679-019-00364-y>)
- 10 Broughton DE & Moley KH. Obesity and female infertility: potential mediators of obesity's impact. *Fertility and Sterility* 2017 **107** 840–847. (<https://doi.org/10.1016/j.fertnstert.2017.01.017>)
- 11 Vrbikova J & Hainer V. Obesity and polycystic ovary syndrome. *Obesity Facts* 2009 **2** 26–35. (<https://doi.org/10.1159/000194971>)
- 12 Estienne A, Bongrani A, Reverchon M, Ramé C, Ducluzeau PH, Froment P & Dupont J. Involvement of novel adipokines, chemerin, visfatin, resistin and apelin in reproductive functions in normal and pathological conditions in humans and animal models. *International Journal of Molecular Sciences* 2019 **20** 4431. (<https://doi.org/10.3390/ijms20184431>)
- 13 Ruiz-Cortés ZT, Martel-Kennes Y, Gévy NY, Downey BR, Palin MF & Murphy BD. Biphasic effects of leptin in porcine granulosa cells. *Biology of Reproduction* 2003 **68** 789–796. (<https://doi.org/10.1095/biolreprod.102.010702>)
- 14 Sirotkin AV, Chrenková M, Nitrayová S, Patraš P, Darlak K, Valenzuela F, Pinilla L & Tena-Sempere M. Effects of chronic food restriction and treatments with leptin or ghrelin on different reproductive parameters of male rats. *Peptides* 2008 **29** 1362–1368. (<https://doi.org/10.1016/j.peptides.2008.03.011>)
- 15 Gregoraszczuk EL, Wójciewicz AK, Ptak A & Nowak K. *In vitro* effect of leptin on steroids' secretion by FSH-and LH-treated porcine small, medium and large preovulatory follicles. *Reproductive Biology* 2003 **3** 227–239.
- 16 Reshma R, Mishra SR, Thakur N, Parmar MS, Somal A, Bharti MK, Pandey S, Chandra V, Chouhan VS, Verma MR, *et al.* Modulatory role of leptin on ovarian functions in water buffalo (*Bubalus bubalis*). *Theriogenology* 2016 **86** 1720–1739. (<https://doi.org/10.1016/j.theriogenology.2016.05.029>)
- 17 Nikanfar S, Oghbaei H, Rezaei YR, Zarezaheh R, Jafari-Gharabaghloou D, Nejabati HR, Bahrami Z, Bleisinger N, Samadi N, Fattahi A, *et al.* Role of adipokines in the ovarian function: oogenesis and steroidogenesis. *Journal of Steroid Biochemistry and Molecular Biology* 2021 **209** 105852. (<https://doi.org/10.1016/j.jsbmb.2021.105852>)
- 18 Rak A, Drwal E, Karpeta A & Gregoraszczuk EL. Regulatory role of gonadotropins and local factors produced by ovarian follicles on *in vitro* resistin expression and action on porcine follicular steroidogenesis. *Biology of Reproduction* 2015 **92** 142. (<https://doi.org/10.1095/biolreprod.115.128611>)
- 19 Rak A, Zajda K & Gregoraszczuk EL. Endocrine disrupting compounds modulates adiponectin secretion, expression of its receptors and action on steroidogenesis in ovarian follicle. *Reproductive Toxicology* 2017 **69** 204–211. (<https://doi.org/10.1016/j.reprotox.2017.03.004>)
- 20 Fernández-Sánchez A, Madrigal-Santillán E, Bautista M, Esquivel-Soto J, Morales-González A, Esquivel-Chirino C, Durante-Montiel I, Sánchez-Rivera G, Valadez-Vega C & Morales-González JA. Inflammation, oxidative stress, and obesity. *International Journal of Molecular Sciences* 2011 **12** 3117–3132. (<https://doi.org/10.3390/ijms12053117>)
- 21 Fontana R & Torre SD. The deep correlation between energy metabolism and reproduction: a view on the effects of nutrition for women fertility. *Nutrients* 2016 **8** 87. (<https://doi.org/10.3390/nu8020087>)
- 22 Roa J & Tena-Sempere M. Connecting metabolism and reproduction: roles of central energy sensors and key molecular mediators. *Molecular and Cellular Endocrinology* 2014 **397** 4–14. (<https://doi.org/10.1016/j.mce.2014.09.027>)
- 23 RoyChoudhury S, Singh A, Gupta NJ, Srivastava S, Joshi MV, Chakravarty B & Chaudhury K. Repeated implantation failure versus repeated implantation success: discrimination at a metabolomic level. *Human Reproduction* 2016 **31** 1265–1274. (<https://doi.org/10.1093/humrep/dew064>)
- 24 Herd JM, Palanisamy SK, Ciller UA, Fan R, Moens P, Smart NA & McFarlane JR. An updated view of leptin on implantation and pregnancy: a review. *Physiological Research* 2014 **63** 543–557. (<https://doi.org/10.33549/physiolres.932674>)
- 25 Gudelska M, Dobrzyn K, Kiezun M, Rytelawska E, Kisielewska K, Kaminska B, Kaminski T & Smolinska N. The expression of chemerin and its receptors (CMKLR1, GPR1, CCRL2) in the porcine uterus during the oestrous cycle and early pregnancy and in trophoblasts and conceptuses. *Animal* 2020 **14** 2116–2128. (<https://doi.org/10.1017/S175173112000097X>)
- 26 Desmarais JA, Cao M, Bateman A & Murphy BD. Spatiotemporal expression pattern of progesterone in embryo implantation and placenta formation suggests a role in cell proliferation, remodeling, and angiogenesis. *Reproduction* 2008 **136** 247–257. (<https://doi.org/10.1530/REP-08-0044>)
- 27 Zheng WL, Sierra-Rivera E, Luan J, Osteen KG & Ong DE. Retinoic acid synthesis and expression of cellular retinol-binding protein and cellular retinoic acid-binding protein type II are concurrent with decidualization of rat uterine stromal cells. *Endocrinology* 2000 **141** 802–808. (<https://doi.org/10.1210/endo.141.2.7323>)
- 28 Kelesidis T, Kelesidis I, Chou S & Mantzoros CS. Narrative review: the role of leptin in human physiology: emerging clinical applications. *Annals of Internal Medicine* 2010 **152** 93–100. (<https://doi.org/10.7326/0003-4819-152-2-201001190-00008>)
- 29 Zhang F, Chen Y, Heiman M & DiMarchi R. Leptin: structure, function and biology. *Vitamins and Hormones* 2005 **71** 345–372. ([https://doi.org/10.1016/S0083-6729\(05\)71012-8](https://doi.org/10.1016/S0083-6729(05)71012-8))
- 30 La Cava A. Leptin in inflammation and autoimmunity. *Cytokine* 2017 **98** 51–58. (<https://doi.org/10.1016/j.cyt.2016.10.011>)
- 31 Trujillo ME, Sullivan S, Harten I, Schneider SH, Greenberg AS & Fried SK. Interleukin-6 regulates human adipose tissue lipid metabolism and leptin production *in vitro*. *Journal of Clinical Endocrinology and Metabolism* 2004 **89** 5577–5582. (<https://doi.org/10.1210/jc.2004-0603>)
- 32 Shalitin S & Phillip M. Role of obesity and leptin in the pubertal process and pubertal growth – a review. *International Journal of Obesity and Related Metabolic Disorders* 2003 **27** 869–874. (<https://doi.org/10.1038/sj.ijo.0802328>)
- 33 Hausman GJ, Barb CR & Lents CA. Leptin and reproductive function. *Biochimie* 2012 **94** 2075–2081. (<https://doi.org/10.1016/j.biochi.2012.02.022>)
- 34 Kitawaki J, Koshihara H, Ishihara H, Kusuki I, Tsukamoto K & Honjo H. Expression of leptin receptor in human endometrium and fluctuation during the menstrual cycle. *Journal of Clinical Endocrinology*

- and Metabolism 2000 **85** 1946–1950. (<https://doi.org/10.1210/jcem.85.5.6567>)
- 35 González RR, Caballero-Campo P, Jasper M, Mercader A, Devoto L, Pellicer A & Simon C. Leptin and leptin receptor are expressed in the human endometrium and endometrial leptin secretion is regulated by the human blastocyst. *Journal of Clinical Endocrinology and Metabolism* 2000 **85** 4883–4888. (<https://doi.org/10.1210/jcem.85.12.7060>)
 - 36 Malik NM, Carter ND, Murray JF, Scaramuzzi RJ, Wilson CA & Stock MJ. Leptin requirement for conception, implantation, and gestation in the mouse. *Endocrinology* 2001 **142** 5198–5202. (<https://doi.org/10.1210/endo.142.12.8535>)
 - 37 Cervero A, Horcajadas JA, Martín J, Pellicer A & Simón C. The leptin system during human endometrial receptivity and preimplantation development. *Journal of Clinical Endocrinology and Metabolism* 2004 **89** 2442–2451. (<https://doi.org/10.1210/jc.2003-032127>)
 - 38 Ramos MP, Rueda BR, Leavis PC & Gonzalez RR. Leptin serves as an upstream activator of an obligatory signaling cascade in the embryo-implantation process. *Endocrinology* 2005 **146** 694–701. (<https://doi.org/10.1210/en.2004-1186>)
 - 39 Tanaka T & Umesaki N. Leptin regulates the proliferation and apoptosis of human endometrial epithelial cells. *International Journal of Molecular Medicine* 2008 **22** 683–689. (<https://doi.org/10.3892/ijmm.00000073>)
 - 40 Yang YJ, Cao YJ, Bo SM, Peng S, Liu WM & Duan EK. Leptin-directed embryo implantation: leptin regulates adhesion and outgrowth of mouse blastocysts and receptivity of endometrial epithelial cells. *Animal Reproduction Science* 2006 **92** 155–167. (<https://doi.org/10.1016/j.anireprosci.2005.05.019>)
 - 41 Tanaka T, Utsunomiya T, Bai T, Nakajima S & Umesaki N. Leptin inhibits decidualization and enhances cell viability of normal human endometrial stromal cells. *International Journal of Molecular Medicine* 2003 **12** 95–98. (<https://doi.org/10.3892/ijmm.12.1.95>)
 - 42 Fukuda J, Nasu K, Sun B, Shang S, Kawano Y & Miyakawa I. Effects of leptin on the production of cytokines by cultured human endometrial stromal and epithelial cells. *Fertility and Sterility* 2003 **80** (Supplement 2) 783–787. ([https://doi.org/10.1016/S0015-0282\(03\)00776-3](https://doi.org/10.1016/S0015-0282(03)00776-3))
 - 43 Dekel N, Gnainsky Y, Granot I, Racicot K & Mor G. The role of inflammation for a successful implantation. *American Journal of Reproductive Immunology* 2014 **72** 141–147. (<https://doi.org/10.1111/aji.12266>)
 - 44 Achari AE & Jain SK. Adiponectin, a therapeutic target for obesity, diabetes, and endothelial dysfunction. *International Journal of Molecular Sciences* 2017 **18** 1321. (<https://doi.org/10.3390/ijms18061321>)
 - 45 Yamauchi T, Iwabu M, Okada-Iwabu M & Kadowaki T. Adiponectin receptors: a review of their structure, function and how they work. *Best Practice and Research: Clinical Endocrinology and Metabolism* 2014 **28** 15–23. (<https://doi.org/10.1016/j.beem.2013.09.003>)
 - 46 Straub LG & Scherer PE. Metabolic messengers: adiponectin. *Nature Metabolism* 2019 **1** 334–339. (<https://doi.org/10.1038/s42255-019-0041-z>)
 - 47 Michalakakis KG & Segars JH. The role of adiponectin in reproduction: from polycystic ovary syndrome to assisted reproduction. *Fertility and Sterility* 2010 **94** 1949–1957. (<https://doi.org/10.1016/j.fertnstert.2010.05.010>)
 - 48 Cheng L, Shi H, Jin Y, Li X, Pan J, Lai Y, Lin Y, Jin Y, Roy G, Zhao A, *et al.* Adiponectin deficiency leads to female subfertility and ovarian dysfunctions in mice. *Endocrinology* 2016 **157** 4875–4887. (<https://doi.org/10.1210/en.2015-2080>)
 - 49 Liu YH, Tsai EM, Chen YL, Chen HS, Chen YC, Wu LC, Lee CH, Jong SB & Chan TF. Serum adiponectin levels increase after human chorionic gonadotropin treatment during *in vitro* fertilization. *Gynecologic and Obstetric Investigation* 2006 **62** 61–65. (<https://doi.org/10.1159/000092260>)
 - 50 Takemura Y, Osuga Y, Yamauchi T, Kobayashi M, Harada M, Hirata T, Morimoto C, Hirota Y, Yoshino O, Koga K, *et al.* Expression of adiponectin receptors and its possible implication in the human endometrium. *Endocrinology* 2006 **147** 3203–3210. (<https://doi.org/10.1210/en.2005-1510>)
 - 51 Smolinska N, Dobrzyn K, Kiezun M, Szeszko K, Maleszka A & Kaminski T. Effect of adiponectin on the steroidogenic acute regulatory protein, P450 side chain cleavage enzyme and 3 β -hydroxysteroid dehydrogenase gene expression, progesterone and androstenedione production by the porcine uterus during early pregnancy. *Journal of Physiology and Pharmacology* 2016 **67** 443–456.
 - 52 Lim W, Choi MJ, Bae H, Bazer FW & Song G. A critical role for adiponectin-mediated development of endometrial luminal epithelial cells during the peri-implantation period of pregnancy. *Journal of Cellular Physiology* 2017 **232** 3146–3157. (<https://doi.org/10.1002/jcp.25768>)
 - 53 Altun T, Jindal S, Greenfield K, Shu J & Pal L. Low follicular fluid IL-6 levels in IVF patients are associated with increased likelihood of clinical pregnancy. *Journal of Assisted Reproduction and Genetics* 2011 **28** 245–251. (<https://doi.org/10.1007/s10815-010-9502-8>)
 - 54 Bohlouli S, Khazaei M, Rabzia A, Khazaei MR, Sadeghi E, Bohlouli S, Khazaei M, Rabzia A, Khazaei M & Sadeghi E. Adiponectin effect on nitric oxide secretion by normal and endometriotic human endometrial stromal cells: *in vitro* study. *International Journal of Morphology* 2015 **33** 337–341. (<https://doi.org/10.4067/S0717-95022015000100053>)
 - 55 Saxena D, Purohit SB, Kumar GP & Laloraya M. Increased appearance of inducible nitric oxide synthase in the uterus and embryo at implantation. *Nitric Oxide: Biology and Chemistry* 2000 **4** 384–391. (<https://doi.org/10.1006/niox.2000.0286>)
 - 56 Kim ST, Marquard K, Stephens S, Loudon E, Allsworth J & Moley KH. Adiponectin and adiponectin receptors in the mouse preimplantation embryo and uterus. *Human Reproduction* 2011 **26** 82–95. (<https://doi.org/10.1093/humrep/deq292>)
 - 57 Kakizawa S. Apelin. In *Handbook of Hormones*, p. 277.e31–33. Amsterdam, Netherlands: Elsevier, 2016. (<https://doi.org/10.1016/B978-0-12-801028-0.00031-3>)
 - 58 Chapman NA, Dupré DJ & Rainey JK. The apelin receptor: physiology, pathology, cell signalling, and ligand modulation of a peptide-activated class A GPCR. *Biochemistry and Cell Biology* 2014 **92** 431–440. (<https://doi.org/10.1139/bcb-2014-0072>)
 - 59 Antushevich H & Wójcik M. Review: Apelin in disease. *Clinica Chimica Acta* 2018 **483** 241–248. (<https://doi.org/10.1016/j.cca.2018.05.012>)
 - 60 Han S, Englander EW, Gomez GA & Greeley Jr GH. Apelin regulates nuclear factor- κ B's involvement in the inflammatory response of pancreatitis. *Pancreas* 2017 **46** 64–70. (<https://doi.org/10.1097/MPA.0000000000000740>)
 - 61 Taheri S, Murphy K, Cohen M, Sujkovic E, Kennedy A, Dhillon W, Dakin C, Sajedi A, Ghatei M & Bloom S. The effects of centrally administered apelin-13 on food intake, water intake and pituitary hormone release in rats. *Biochemical and Biophysical Research Communications* 2002 **291** 1208–1212. (<https://doi.org/10.1006/bbrc.2002.6575>)
 - 62 Rak A, Drwal E, Rame C, Knapczyk-Stwora K, Słomczyńska M, Dupont J & Gregoraszczuk EL. Expression of apelin and apelin receptor (APJ) in porcine ovarian follicles and *in vitro* effect of apelin on steroidogenesis and proliferation through APJ activation and different signaling pathways. *Theriogenology* 2017 **96** 126–135. (<https://doi.org/10.1016/j.theriogenology.2017.04.014>)
 - 63 Ozkan ZS, Cilgin H, Simsek M, Cobanoglu B & Ilhan N. Investigation of apelin expression in endometriosis. *Journal of Reproduction and Infertility* 2013 **14** 50–55.
 - 64 Różycka M, Kurowska P, Grzesiak M, Kotula-Balak M, Tworzydło W, Rame C, Gregoraszczuk E, Dupont J & Rak A. Apelin and apelin receptor at different stages of corpus luteum development and effect of apelin on progesterone secretion and 3 β -hydroxysteroid dehydrogenase (3 β -HSD) in pigs. *Animal Reproduction Science* 2018 **192** 251–260. (<https://doi.org/10.1016/j.anireprosci.2018.03.021>)
 - 65 Schilffarth S, Antoni B, Schams D, Meyer HH & Berisha B. The expression of apelin and its receptor APJ during different physiological

- stages in the bovine ovary. *International Journal of Biological Sciences* 2009 **5** 344–350. (<https://doi.org/10.7150/ijbs.5.344>)
- 66 Xu F & Stouffer RL. Dynamic expression of apelin and its receptor in the primate preovulatory follicle and corpus luteum during the menstrual cycle. *Biology of Reproduction* 2012 **87** Suppl_1 169. (<https://doi.org/10.1093/biolreprod/87.s1.169>)
 - 67 Mlyczyńska E, Kurowska P, Drwal E, OpydoChanek M, Tworzydło W, Kotula-Balak M & Rak A. Apelin and apelin receptor in human placenta: expression, signalling pathway and regulation of trophoblast JEG3 and BeWo cells proliferation and cell cycle. *International Journal of Molecular Medicine* 2020 **45** 691–702. (<https://doi.org/10.3892/ijmm.2020.4452>)
 - 68 Griffiths RM, Pru CA, Behura SK, Cronrath AR, McCallum ML, Kelp NC, Winuthayanon W, Spencer TE & Pru JK. AMPK is required for uterine receptivity and normal responses to steroid hormones. *Reproduction* 2020 **159** 707–717. (<https://doi.org/10.1530/REP-19-0402>)
 - 69 Hehir MP & Morrison JJ. The adipokine apelin and human uterine contractility. *American Journal of Obstetrics and Gynecology* 2012 **206** 359.e1–359.e5. (<https://doi.org/10.1016/j.ajog.2012.01.032>)
 - 70 Kacar E, Ercan Z, Serhatlioglu I, Sumer A, Kelestimur H & Kutlu S. The effects of apelin on myometrium contractions in pregnant rats. *Cellular and Molecular Biology* 2018 **64** 74–79. (<https://doi.org/10.14715/cmb/2018.64.11.13>)
 - 71 Foussal C, Lairez O, Calise D, Pathak A, Guilbeau-Frugier C, Valet P, Parini A & Kunduzova O. Activation of catalase by apelin prevents oxidative stress-linked cardiac hypertrophy. *FEBS Letters* 2010 **584** 2363–2370. (<https://doi.org/10.1016/j.febslet.2010.04.025>)
 - 72 Than A, Zhang X, Leow MK-S, Poh CL, Chong SK & Chen P. Apelin attenuates oxidative stress in human adipocytes. *Journal of Biological Chemistry* 2014 **289** 3763–3774. (<https://doi.org/10.1074/jbc.M113.526210>)
 - 73 Guerin P, El Mouatassim S & Menezo Y. Oxidative stress and protection against reactive oxygen species in the pre-implantation embryo and its surroundings. *Human Reproduction Update* 2001 **7** 175–189. (<https://doi.org/10.1093/humupd/7.2.175>)
 - 74 Shahin S, Singh VP, Shukla RK, Dhawan A, Gangwar RK, Singh SP & Chaturvedi CM. 2.45 GHz microwave irradiation-induced oxidative stress affects implantation or pregnancy in mice, *Mus musculus*. *Applied Biochemistry and Biotechnology* 2013 **169** 1727–1751. (<https://doi.org/10.1007/s12010-012-0079-9>)
 - 75 Hu J, Cheng D, Gao X, Bao J, Ma X & Wang H. Vitamin C enhances the *in vitro* development of porcine pre-implantation embryos by reducing oxidative stress. *Reproduction in Domestic Animals* 2012 **47** 873–879. (<https://doi.org/10.1111/j.1439-0531.2011.01982.x>)
 - 76 Helfer G & Wu QF. Chemerin: a multifaceted adipokine involved in metabolic disorders. *Journal of Endocrinology* 2018 **238** R79–R94. (<https://doi.org/10.1530/JOE-18-0174>)
 - 77 Kaur J, Adya R, Tan BK, Chen J & Randeve HS. Identification of chemerin receptor (ChemR23) in human endothelial cells: chemerin-induced endothelial angiogenesis. *Biochemical and Biophysical Research Communications* 2010 **391** 1762–1768. (<https://doi.org/10.1016/j.bbrc.2009.12.150>)
 - 78 Kim JY, Xue K, Cao M, Wang Q, Liu JY, Leader A, Han JY & Tsang BK. Chemerin suppresses ovarian follicular development and its potential involvement in follicular arrest in rats treated chronically with dihydrotestosterone. *Endocrinology* 2013 **154** 2912–2923. (<https://doi.org/10.1210/en.2013-1001>)
 - 79 Wang Q, Kim JY, Xue K, Liu JY, Leader A & Tsang BK. Chemerin, a novel regulator of follicular steroidogenesis and its potential involvement in polycystic ovarian syndrome. *Endocrinology* 2012 **153** 5600–5611. (<https://doi.org/10.1210/en.2012-1424>)
 - 80 Dobrzyn K, Kiezun M, Zaobidna E, Kisieleska K, Rytelska E, Gudelska M, Kopij G, Bors K, Szymanska K, Kaminska B, *et al.* The *in vitro* effect of prostaglandin E2 and F2 α on the chemerin system in the porcine endometrium during gestation. *International Journal of Molecular Sciences* 2020 **21** 5213. (<https://doi.org/10.3390/ijms21155213>)
 - 81 Yang X, Yao J, Wei Q, Ye J, Yin X, Quan X, Lan Y & Xing H. Role of chemerin/CMKLR1 in the maintenance of early pregnancy. *Frontiers of Medicine* 2018 **12** 525–532. (<https://doi.org/10.1007/s11684-017-0577-9>)
 - 82 Shen W, Tian C, Chen H, Yang Y, Zhu D, Gao P & Liu J. Oxidative stress mediates chemerin-induced autophagy in endothelial cells. *Free Radical Biology and Medicine* 2013 **55** 73–82. (<https://doi.org/10.1016/j.freeradbiomed.2012.11.011>)
 - 83 Carlino C, Trotta E, Stabile H, Morrone S, Bulla R, Soriani A, Iannitto ML, Agostinis C, Mocci C, Minozzi M, *et al.* Chemerin regulates NK cell accumulation and endothelial cell morphogenesis in the decidua during early pregnancy. *Journal of Clinical Endocrinology and Metabolism* 2012 **97** 3603–3612. (<https://doi.org/10.1210/jc.2012-1102>)
 - 84 Sojka DK, Yang L & Yokoyama WM. Uterine natural killer cells. *Frontiers in Immunology* 2019 **10** 960. (<https://doi.org/10.3389/fimmu.2019.00960>)
 - 85 Boomsma CM, Kavelaars A, Eijkemans MJ, Fauser BC, Heijnen CJ & Macklon NS. Ovarian stimulation for *in vitro* fertilization alters the intrauterine cytokine, chemokine, and growth factor milieu encountered by the embryo. *Fertility and Sterility* 2010 **94** 1764–1768. (<https://doi.org/10.1016/j.fertnstert.2009.10.044>)
 - 86 Mariani F & Roncucci L. Chemerin/chemR23 axis in inflammation onset and resolution. *Inflammation Research* 2015 **64** 85–95. (<https://doi.org/10.1007/s00011-014-0792-7>)
 - 87 Parolini S, Santoro A, Marcenaro E, Luini W, Massardi L, Facchetti F, Communi D, Parmentier M, Majorana A, Sironi M, *et al.* The role of chemerin in the colocalization of NK and dendritic cell subsets into inflamed tissues. *Blood* 2007 **109** 3625–3632. (<https://doi.org/10.1182/blood-2006-08-038844>)
 - 88 Yu S, Zhang Y, Li MZ, Xu H, Wang Q, Song J, Lin P, Zhang L, Liu Q, Huang QX, *et al.* Chemerin and apelin are positively correlated with inflammation in obese type 2 diabetic patients. *Chinese Medical Journal* 2012 **125** 3440–3444. (<https://doi.org/10.3760/cma.j.issn.0366-6999.2012.19.015>)
 - 89 Gnainsky Y, Granot I, Aldo PB, Barash A, Or Y, Schechtman E, Mor G & Dekel N. Local injury of the endometrium induces an inflammatory response that promotes successful implantation. *Fertility and Sterility* 2010 **94** 2030–2036. (<https://doi.org/10.1016/j.fertnstert.2010.02.022>)
 - 90 Yamawaki H, Kameshima S, Usui T, Okada M & Hara Y. A novel adipocytokine, chemerin exerts anti-inflammatory roles in human vascular endothelial cells. *Biochemical and Biophysical Research Communications* 2012 **423** 152–157. (<https://doi.org/10.1016/j.bbrc.2012.05.103>)
 - 91 Adeghate E. Visfatin: structure, function and relation to diabetes mellitus and other dysfunctions. *Current Medicinal Chemistry* 2008 **15** 1851–1862. (<https://doi.org/10.2174/092986708785133004>)
 - 92 Sommer G, Garten A, Petzold S, Beck-Sickinger AG, Blüher M, Stumvoll M & Fasshauer M. Visfatin/PBEF/Nampt: structure, regulation and potential function of a novel adipokine. *Clinical Science* 2008 **115** 13–23. (<https://doi.org/10.1042/CS20070226>)
 - 93 Peiró C, Romacho T, Carraro R & Sánchez-Ferrer CE. Visfatin/PBEF/Nampt: a new cardiovascular target? *Frontiers in Pharmacology* 2010 **1** 135. (<https://doi.org/10.3389/fphar.2010.00135>)
 - 94 Tilg H & Moschen AR. Role of adiponectin and PBEF/visfatin as regulators of inflammation: involvement in obesity-associated diseases. *Clinical Science* 2008 **114** 275–288. (<https://doi.org/10.1042/CS20070196>)
 - 95 Camp SM, Ceco E, Evenoski CL, Danilov SM, Zhou T, Chiang ET, Moreno-Vinasco L, Mapes B, Zhao J, Gursoy G, *et al.* Unique toll-like receptor 4 activation by NAMPT/PBEF induces NF κ B signaling and inflammatory lung injury. *Scientific Reports* 2015 **5** 13135. (<https://doi.org/10.1038/srep13135>)
 - 96 Fukuhara A, Matsuda M, Nishizawa M, Segawa K, Tanaka M, Kishimoto K, Matsuki Y, Murakami M, Ichisaka T, Murakami H, *et al.* Visfatin: a protein secreted by visceral fat that mimics the effects of insulin. *Science* 2005 **307** 426–430. (<https://doi.org/10.1126/science.1097243>)

- 97 Yang Z, Xiao K, Wang W, Tang J, Sun P-P, Peng K-M & Song H. The effect of visfatin on inflammatory reaction in uterus of LPS-induced rats. *International Journal of Morphology* 2015 **33** 194–203. (<https://doi.org/10.4067/S0717-95022015000100031>)
- 98 Wyskida K, Franik G, Wikarek T, Owczarek A, Delroba A, Chudek J, Sikora J & Olszanecka-Glinianowicz M. The levels of adipokines in relation to hormonal changes during the menstrual cycle in young, normal-weight women. *Endocrine Connections* 2017 **6** 892–900. (<https://doi.org/10.1530/EC-17-0186>)
- 99 Annie L, Gurusubramanian G & Roy VK. Estrogen and progesterone dependent expression of visfatin/NAMPT regulates proliferation and apoptosis in mice uterus during estrous cycle. *Journal of Steroid Biochemistry and Molecular Biology* 2019 **185** 225–236. (<https://doi.org/10.1016/j.jsbmb.2018.09.010>)
- 100 Kim SC, Park MJ, Joo BS, Joo JK, Suh DS & Lee KS. Decreased expressions of vascular endothelial growth factor and visfatin in the placental bed of pregnancies complicated by preeclampsia. *Journal of Obstetrics and Gynaecology Research* 2012 **38** 665–673. (<https://doi.org/10.1111/j.1447-0756.2011.01773.x>)
- 101 Mastorakos G, Valsamakis G, Papatheodorou DC, Barlas I, Margeli A, Boutsiadis A, Kouskouni E, Vitoratos N, Papadimitriou A, Papassotiropoulos I, *et al.* The role of adipocytokines in insulin resistance in normal pregnancy: visfatin concentrations in early pregnancy predict insulin sensitivity. *Clinical Chemistry* 2007 **53** 1477–1483. (<https://doi.org/10.1373/clinchem.2006.084731>)
- 102 Huang Z, Liu Y, Qi G, Brand D & Zheng SG. Role of vitamin A in the immune system. *Journal of Clinical Medicine* 2018 **7** 258. (<https://doi.org/10.3390/jcm7090258>)
- 103 Clagett-Dame M & Knutson D. Vitamin A in reproduction and development. *Nutrients* 2011 **3** 385–428. (<https://doi.org/10.3390/nu3040385>)
- 104 Perduca M, Nicolis S, Mannucci B, Galliano M & Monaco HL. Human plasma retinol-binding protein (RBP4) is also a fatty acid-binding protein. *Biochimica et Biophysica Acta: Molecular and Cell Biology of Lipids* 2018 **1863** 458–466. (<https://doi.org/10.1016/j.bbalip.2018.01.010>)
- 105 Li H, Cao G, Zhang N, Lou T, Wang Q, Zhang Z & Liu C. RBP4 regulates trophoblastic cell proliferation and invasion via the PI3K/AKT signaling pathway. *Molecular Medicine Reports* 2018 **18** 2873–2879. (<https://doi.org/10.3892/mmr.2018.9240>)
- 106 Dancis J, Levitz M, Katz J, Wilson D, Blaser WS, Piantadosi R & Goodman DS. Transfer and metabolism of retinol by the perfused human placenta. *Pediatric Research* 1992 **32** 195–199. (<https://doi.org/10.1203/00006450-199208000-00014>)
- 107 Pavone ME, Malpani S, Dyson M & Bulun SE. Altered retinoid signaling compromises decidualization in human endometrial stromal cells. *Reproduction* 2017 **154** 207–216. (<https://doi.org/10.1530/REP-16-0592>)
- 108 Mullen MP, Forde N, Parr MH, Diskin MG, Morris DG, Nally JE, Evans AC & Crowe MA. Alterations in systemic concentrations of progesterone during the early luteal phase affect RBP4 expression in the bovine uterus. *Reproduction, Fertility, and Development* 2012 **24** 715–722. (<https://doi.org/10.1071/RD11246>)
- 109 Dassen H, Punyadeera C, Kamps R, Klomp J, Dunselman G, Dijkjes F, De Goeij A, Ederveen A & Groothuis P. Progesterone regulation of implantation-related genes: new insights into the role of oestrogen. *Cellular and Molecular Life Sciences* 2007 **64** 1009–1032. (<https://doi.org/10.1007/s00018-007-6553-9>)
- 110 Chan TF, Chen HS, Chen YC, Lee CH, Chou FH, Chen IJ, Chen SY, Jong SB & Tsai EM. Increased serum retinol-binding protein 4 concentrations in women with gestational diabetes mellitus. *Reproductive Sciences* 2007 **14** 169–174. (<https://doi.org/10.1177/1933719106298407>)
- 111 Cheng YH, Yin P, Xue Q, Yilmaz B, Dawson MI & Bulun SE. Retinoic acid (RA) regulates 17 β -hydroxysteroid dehydrogenase type 2 expression in endometrium: interaction of RA receptors with specificity protein (SP) 1/SP3 for estradiol metabolism. *Journal of Clinical Endocrinology and Metabolism* 2008 **93** 1915–1923. (<https://doi.org/10.1210/jc.2007-1536>)
- 112 Bulun SE, Cheng YH, Pavone ME, Yin P, Imir G, Utsunomiya H, Thung S, Xue Q, Marsh EE & Tokunaga H. 17 β -Hydroxysteroid dehydrogenase-2 deficiency and progesterone resistance in endometriosis. *Seminars in Reproductive Medicine* 2010 **28** 44–50. (<https://doi.org/10.1055/s-0029-1242992>)
- 113 Forde N & Loneragan P. Transcriptomic analysis of the bovine endometrium: what is required to establish uterine receptivity to implantation in cattle? *Journal of Reproduction and Development* 2012 **58** 189–195. (<https://doi.org/10.1262/jrd.2011-021>)
- 114 Toh H, Chitramuthu BP, Bennett HP & Bateman A. Structure, function, and mechanism of progranulin; the brain and beyond. *Journal of Molecular Neuroscience* 2011 **45** 538–548. (<https://doi.org/10.1007/s12031-011-9569-4>)
- 115 Pogonowska M, Poniatowski LA, Wawrzyniak A, Królikowska K & Kalicki B. The role of progranulin (PGRN) in the modulation of anti-inflammatory response in asthma. *Central-European Journal of Immunology* 2019 **44** 97–101. (<https://doi.org/10.5114/ceji.2019.83267>)
- 116 Tanaka K, Osaka M, Takemori S, Watanabe M, Tanigaki S & Kobayashi Y. Contraction-associated proteins expression by human uterine smooth muscle cells depends on maternal serum and progranulin associated with gestational weight gain. *Endocrine Journal* 2020 **67** 819–825. (<https://doi.org/10.1507/endocrj.EJ20-0037>)
- 117 Diaz-Cueto L, Stein P, Jacobs A, Schultz RM & Gerton GL. Modulation of mouse preimplantation embryo development by acrogranin (epithelin/granulin precursor). *Developmental Biology* 2000 **217** 406–418. (<https://doi.org/10.1006/dbio.1999.9564>)
- 118 Qin J, Diaz-Cueto L, Schwarze JE, Takahashi Y, Imai M, Isuzugawa K, Yamamoto S, Chang KT, Gerton GL & Imakawa K. Effects of progranulin on blastocyst hatching and subsequent adhesion and outgrowth in the mouse. *Biology of Reproduction* 2005 **73** 434–442. (<https://doi.org/10.1095/biolreprod.105.040030>)
- 119 Daniel R, Daniels E, He Z & Bateman A. Progranulin (acrogranin/PC cell-derived growth factor/granulin-epithelin precursor) is expressed in the placenta, epidermis, microvasculature, and brain during murine development. *Developmental Dynamics* 2003 **227** 593–599. (<https://doi.org/10.1002/dvdy.10341>)
- 120 Perez CM, Minimo C, Caballero-Campo P, Rinaudo P, Orris J & Gerton GL. P-537: progranulin characterization in human endometrium: translational studies from mouse to human. *Fertility and Sterility* 2006 **86** S333. (<https://doi.org/10.1016/j.fertnstert.2006.07.906>)
- 121 Perez CM, Diaz-cueto L, Van der voort D, Schwarze JE & Gerton GL. Acrogranin (granulin epithelin precursor, PCDFG) is crucial for mouse embryo receptivity and implantation. *Fertility and Sterility* 2003 **80** 27–28. ([https://doi.org/10.1016/S0015-0282\(03\)01873-9](https://doi.org/10.1016/S0015-0282(03)01873-9))
- 122 Xu B, Chen X, Ding Y, Chen C, Liu T & Zhang H. Abnormal angiogenesis of placenta in progranulin-deficient mice. *Molecular Medicine Reports* 2020 **22** 3482–3492. (<https://doi.org/10.3892/mmr.2020.11438>)
- 123 Gonzalez RR, Devoto L, Campana A & Bischof P. Effects of leptin, interleukin-1 α , interleukin-6, and transforming growth factor-beta on markers of trophoblast invasive phenotype: integrins and metalloproteinases. *Endocrine* 2001 **15** 157–164. (<https://doi.org/10.1385/endo.15.2.157>)
- 124 Schulz LC & Widmaier EP. The effect of leptin on mouse trophoblast cell invasion. *Biology of Reproduction* 2004 **71** 1963–1967. (<https://doi.org/10.1095/biolreprod.104.032722>)

Received in final form 24 August 2021

Accepted 24 September 2021

Accepted Manuscript published online 24 September 2021