ERRATUM

Adipokine expression and endothelial function in subclinical hypothyroidism rats

Ningning Gong^{1,2,*}, Cuixia Gao^{3,*}, Xuedi Chen¹, Yu Wang¹ and Limin Tian¹

¹Department of Endocrinology, Gansu Provincial Hospital, Lanzhou, China

²Department of Clinical Medicine, Gansu University of Chinese Medicine, Lanzhou, Gansu, China

³Department of Ultrasonic Diagnosis, Gansu Provincial Hospital, Lanzhou, China

Correspondence should be addressed to L Tian: tlim6666@sina.com

*(N Gong and C Gao contributed equally to this work)

The authors and journal apologise for an error in the above paper, which appeared in volume 7 part 2, pages 295–304. The error relates to the text in the ‘Animal model’ section of the ‘Materials and methods’ section given on page 296:

The original article stated:

‘In the sHT rats, groups A–C were separately induced by administration of 20 mg/kg/day methimazole (MMI), 30 mg/kg/day MMI and 40 mg/kg/day MMI once daily by gavage.’

This should have stated:

‘In the sHT rats, groups A–C were separately induced by administration of 5 mg/kg/day methimazole (MMI), 15 mg/kg/day MMI and 20 mg/kg/day MMI once daily by gavage.’