

## REVIEW

# Potential genetic polymorphisms predicting polycystic ovary syndrome

Yao Chen and Shu-ying Fang

Hangzhou Fuyang Women and Children Hospital, Hangzhou, China

Correspondence should be addressed to Y Chen: [fychy@163.com](mailto:fychy@163.com)

## Abstract

Polycystic ovary syndrome (PCOS) is a heterogenous endocrine disorder with typical symptoms of oligomenorrhoea, hyperandrogenism, hirsutism, obesity, insulin resistance and increased risk of type 2 diabetes mellitus. Extensive evidence indicates that PCOS is a genetic disease and numerous biochemical pathways have been linked with its pathogenesis. A number of genes from these pathways have been investigated, which include those involved with steroid hormone biosynthesis and metabolism, action of gonadotropin and gonadal hormones, folliculogenesis, obesity and energy regulation, insulin secretion and action and many others. In this review, we summarize the historical and recent findings in genetic polymorphisms of PCOS from the relevant publications and outline some genetic polymorphisms that are potentially associated with the risk of PCOS. This information could uncover candidate genes associating with PCOS, which will be valuable for the development of novel diagnostic and treatment platforms for PCOS patients.

### Key Words

- ▶ polycystic ovary syndrome
- ▶ genetic polymorphisms
- ▶ candidate gene
- ▶ endocrine

Endocrine Connections  
(2018) 7, R187–R195

## Introduction

Polycystic ovary syndrome (PCOS) is a heterogeneous and complex disorder of reproductive aged women, which usually causes reproduction and metabolism failure (1, 2). It often presents as hyperandrogenism, hyperinsulinemia, insulin resistance, ovulatory dysfunction and polycystic ovaries. The prevalence of PCOS may be as high as 15–20% of women of reproductive age (3). Although it was first reported in 1935 by Stein *et al.*, its etiology is still unclear. PCOS has a complicated and ambiguous pathogenesis. Many risk factors have been identified, such as obesity, genetics and early childhood growth patterns (4, 5).

Data from many studies suggest that genetics is very important in the development of PCOS. A Dutch twin study supported the genetic contribution, with PCOS heritability predicted to be as high as 0.79 (6). Family studies also clarified the effects of genetics on PCOS (7, 8, 9) and its concomitant representations such as hyperandrogenemia (10) and insulin resistance (11). Current studies of the genetics of PCOS mainly

concentrate on multiple candidate genes, benefiting from the advances in technology including the completed HapMap project and genome-wide scans. Candidate genes are usually chosen according to the available information for the potential pathophysiology of the symptoms. These genes are tested by analysis of the affected individuals (12). Several candidate genes in PCOS development have been clarified, including genes associated with steroid biosynthesis and action, gonadotropic secretion and action, folliculogenesis, weight and energy regulation and insulin action (13).

In this review, we aimed to provide an overview of the main genetic polymorphisms that were found or investigated in PCOS patients until now. Relevant literatures of case-control researches and meta-analysis studies were searched from PubMed database or by citation search and reference lists of relevant articles. Relationships of the candidate genes with PCOS that were found or discussed in these publications were summarized.



The genetic polymorphisms are classified as steroidogenic acute regulatory (*StAR*) gene polymorphisms, gonadotropin-releasing hormone receptor (*GnRHR*) gene polymorphisms, follicle-stimulating hormone receptor (*FSHR*) gene polymorphisms, fat mass and obesity-associated (*FTO*) gene polymorphisms, insulin receptor (*IR*) and IR substrate (*IRS*) polymorphisms, vitamin D receptor (*VDR*) polymorphisms, methylenetetrahydrofolate reductase (*MTHFR*) C677T polymorphism, peroxisome proliferator-activated receptor gamma (*PPAR-γ*) polymorphism and interleukin-1beta/IL-1 receptor antagonist (*IL-1β/IL-1Ra*) gene polymorphisms.

### **StAR polymorphisms**

Hyperandrogenism accounts for most of the etiology of PCOS, and it can disturb the steroidogenesis process. The StAR protein regulates the transport of cholesterol through the mitochondrial membrane in the first step of androgen biosynthesis (14, 15). Evidence suggests that patients with PCOS may have an abnormality in androgen biosynthesis, with StAR as one possible target of abnormality (16). Kashar-Miller *et al.* (16) hypothesized that increased production or concentration of StAR may be responsible for the increased ovarian and adrenal androgen found in PCOS of some patients. The authors provided evidence that StAR might participate in the earliest steps of androgen biosynthesis in PCOS. Data from this and subsequent investigations have identified the StAR gene as a candidate gene for PCOS (17). However, the association between *StAR* genetic polymorphisms with the pathology of PCOS remains unclear. In a case-control study, seven *StAR* gene polymorphisms have been detected using polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) method in 45 PCOS women and 80 controls. No relationships between most single nucleotide polymorphisms (SNPs) of the *StAR* gene with PCOS were evident (18). The authors suggested that further evaluation of changes in amino acid active sites and their effect on steroid biosynthesis will be helpful to uncover the association of *StAR* with PCOS. Use of larger study populations would also help definitively identify the exact roles of *StAR* genetic polymorphisms in PCOS.

### **GnRHR polymorphisms**

PCOS patients often have increased levels of plasma luteinizing hormone (LH) and lower levels of follicle-stimulating hormone (FSH) compared to the normal range. These changes might be due to the

dysfunction of the GnRH pulse generator that displays decreased sensitivity to the inhibition of ovarian steroids (19). GnRHR is a G-protein-coupled receptor (GPCR) that is mainly expressed in the gonadotrope membrane of the anterior pituitary (20, 21). GnRH and its receptor are expressed in many extrapituitary tissues, including ovary, placenta, breast and cancer tissues (22). However, the effects of GnRHR on these extrapituitary tissues, especially the ovary, remain unclear. A study by Chen *et al.* involved 314 PCOS patients and 320 control individuals, in which *GnRHR* gene polymorphisms were detected using the PCR-RFLP assay. The results revealed that haplotypes with TCC, CCC and CCT increased the risk of PCOS, while haplotypes with TTT, TCT and TTC lowered the risk (23). A recent genome-wide analysis by whole-exome sequencing and Sanger sequencing confirmation studied three sisters of a consanguineous family diagnosed with PCOS. The results suggested that genetic variations in the hypothalamo-pituitary axis might contribute to the pathogenesis of PCOS (24).

### **FSHR polymorphisms**

FSH participates in the processes of steroidogenesis, oocyte maturation and follicle development via the FSHR located on the surface of granulosa cells in ovary tissue (25). The FSHR is a G-protein-coupled receptor comprising transmembrane, intracellular and extracellular domains. The *FSHR* gene, which is located on chromosome 2p21, has 10 exons and 9 introns (25). Only two *FSHR* polymorphisms are known (p.Thr307Ala and p.Asn680Ser); both are located in exon 10 (17, 26, 27). The p.Asn680Ser variant allele is associated with a higher basal FSH level, longer menstrual cycles and higher ovarian threshold to ovulation induction, reflecting the difference of receptor sensitivity (28, 29, 30, 31, 32). Based on the knowledge that PCOS is characterized by follicular growth failure, several studies broadly investigated the association between p.Thr307Ala and p.Asn680Ser for *FSHR* genetic variations with PCOS. The findings were often inconsistent and controversial. Jin *et al.* (25) performed genotyping in 377 women with PCOS and 388 women without PCOS by TaqMan allelic discrimination assay and found that both p.Thr307Ala and p.Asn680Ser had a significant association with PCOS. The variant homozygote genotypes could increase the risk of PCOS. However, another two studies that used the same genotyping method did not confirm these associations (33, 34). Further studies are needed to better understand the effect of *FSHR* gene polymorphisms in PCOS.

## FTO polymorphisms

The human *FTO* gene is located on chromosome 16q12.2. The gene product is a 2-oxoglutarate-dependent nucleic acid demethylase that is expressed in almost all tissues (35, 36), where it participates in energy metabolism (37). A genome-wide association study in 2007 associated the *FTO* gene with body mass index (BMI) and obesity (38). Obesity is a common characteristic of PCOS; more than 50% of PCOS patients are overweight (39). Thus, it is reasonable to suppose that the *FTO* gene might take part in the pathogenesis of PCOS through BMI and/or obesity. This supposition needs to be investigated. Recent studies identified a common SNP (rs9939609) in the first intron of the *FTO* gene with a T-to-A change in PCOS cases. However, whether *FTO* gene polymorphisms participate in PCOS is still controversial (40, 41, 42, 43, 44, 45, 46). A recent meta-analysis indicated that the *FTO* rs9939609 polymorphism is related to the risk of PCOS in East Asians rather than the overall population (47). Another meta-analysis that included five studies comprising 5010 PCOS cases and 5300 controls indicated the relevance of the rs9939609 A/T polymorphism of the *FTO* gene with PCOS risk and that an allele might be a risk factor for pathology of PCOS (48). Further studies involving more people or the consideration of distinct ethnic and geographical groups will help uncover the effects of *FTO* gene polymorphism on PCOS development.

## IR and IRS polymorphisms

Insulin resistance is an important factor for etiology of PCOS, with a decreased response of glucose to insulin present in over 40% of PCOS cases (49). Genes related to insulin resistance have been widely investigated as risk factors of PCOS. IR and insulin receptor substrate (IRS) proteins are important role in insulin signal transduction. Many studies have explored the effects of *IR* and *IRS* gene polymorphisms on the etiology of PCOS, with a focus on His 1058 C/T for *IR* polymorphism, Gly972Arg for *IRS-1* polymorphism and Gly1057Asp for *IRS-2* polymorphism (50). The results have largely been inconsistent, and no single polymorphism has been directly associated with PCOS. Shi *et al.* (50) recently conducted a meta-analysis to evaluate the relevance between *IR* and *IRS* gene polymorphisms with the pathology of PCOS. The analysis of 28 articles comprising 2975 PCOS cases and 3011 controls suggested an association between the Gly972Arg polymorphism of *IRS-1* and PCOS in Caucasians and between the Gly1057Asp polymorphism of *IRS-2* and

PCOS in Asians. However, no association of the His 1058 C/T polymorphism for *IR* and PCOS was evident.

## VDR polymorphisms

VDR is a DNA-binding transcription factor that is widely distributed in almost all tissues (51). When vitamin D binds to the retinoid X receptor, it enhances the interplay between VDR and this receptor, resulting in the transcription of several vitamin D-regulated genes (52). Vitamin D has a great effect in the metabolic pathways influenced by PCOS (53), and the level of vitamin D is associated with the severity of PCOS. Moreover, *VDR* polymorphisms are related with some characteristics of PCOS. Several studies have investigated the possible relationships of these polymorphisms with the PCOS phenotype mostly using PCR-RFLP method. Mahmoudi *et al.* (54) studied the frequency of FokI, BsmI, ApaI and TaqI *VDR* polymorphisms in healthy Iranian women and Iranian women with PCOS. The CC genotype of ApaI was accompanied by a higher risk of PCOS. Vakili *et al.* (55) demonstrated the dominance of the GG genotype in mild PCOS. GA/AA and A have been linked with the progression of PCOS. El-Shal *et al.* (56) confirmed that the higher frequencies of the C and CC genotypes of TaqI in PCOS cases. However, Ranjzad *et al.* (57) did not observe a statistical difference in ApaI between the women with PCOS and controls. As analyzed by PCR and minisequencing, Jedrzejuk *et al.* (58) found no significantly difference between PCOS and healthy patients concerning the distribution of those polymorphisms. The different data could be explained by the ethnic factor and numbers of classic PCOS women, which need further studies.

## MTHFR C677T polymorphism

MTHFR is important in the processing of 5,10-methylenetetrahydrofolate to 5-methyltetrahydrofolate, which is required for the transformation of homocysteine to methionine (59, 60, 61). An abnormal level of folates inhibits ovulation in immature superovulated rats (62). Folate-deficient monkeys have degraded Graffian follicles and an increased number of atretic and cystic follicles (63). In humans, clinical evidence indicates that PCOS patients usually have an abnormal homocysteine level, which can return to the normal range with folic acid supplementation (64, 65, 66). The collective data implicate *MTHFR* as an ideal candidate gene for PCOS.

**Table 1** Association of gene polymorphisms with the risk of PCOS.

Gene	Description	Polymorphism	Population	Number of PCOS patients	Methods	Association	Ref.
StAR	Steroidogenic acute regulatory	rs104894086	Iranian	45	PCR-RFLP	N	(18)
		rs104894089	Iranian	45		N	(18)
		rs104894090	Iranian	45		N	(18)
		rs137852689	Iranian	45		N	(18)
		rs10489487	Iranian	45		N	(18)
GnRHR	Gonadotropin-releasing hormone receptor	rs104894085	Iranian	45	PCR-RFLP	N	(18)
		rs12644822	Chinese	314		Y	(23)
		rs3756159	Chinese	314		Y	(23)
		rs13138607	Chinese	314		Y	(23)
FSHR	Follicle-stimulating hormone receptor	rs104893836	French	3	Whole-exome sequencing	Y	(24)
		rs6165	Korean	377		TADA	Y
FTO	Fat mass and obesity associated gene	rs6166	Caucasian, Chinese	518 (33), 384 (34)	TADA, sequencing analysis	N	(33, 34)
			Korean, Caucasian, Chinese	377	TADA	Y	(25)
		rs9939609	Chinese, European, Korean	215 (40), 463 (41), 3599 (42), 552 (44)	Sequencing analysis, TaqMan assay, GWAS	Y	(40, 41, 42, 44)
IR	Insulin receptor	rs8050136	Brazilian, American	199 (43), 395 (46)	TADA	N	(43, 46)
			Brazilian, American	199 (43), 395 (46)		N	(43, 46)
		rs1421085	American	395		N	(46)
		rs17817449	American	395		N	(46)
		rs9930506	American	395		N	(46)
IRS	Insulin receptor substrate	rs1799817	Chinese, Turk	2975	Meta-analysis	Y	(50)
			UK, Korean, Japanese, Indian, Iranian, Iraqi, Croatian	2975		N	(50)
VDR	Vitamin D receptor	rs1801278	Greek	2975	Meta-analysis	Y	(50)
			French, Chilean, German, Japanese, American, Spanish, Turk, Chinese, Italian, Indian, Greek, Slovakian, Croatian	2975		N	(50)
VDR	Vitamin D receptor	rs1805097	French, Chinese	2975	PCR-RFLP	Y	(54)
			German, Spanish, Greek	2975		N	(56, 58)
		rs7975232	Iranian	162	PCR-RFLP, minisequencing	Y	(54, 58)
		Egyptian, Polish	150 (56), 92 (58)	N		(54, 58)	
		rs10735810	Iranian, Polish	162 (54), 92 (58)	PCR-RFLP	N	(54, 58)
rs1544410	Iranian, Polish	162 (54), 92 (58)	Y	(56)			
rs731236	Egyptian	150	PCR-RFLP, minisequencing	N	(54, 58)		
Iranian, Polish	162 (54), 92 (58)	N		(54, 58)			
rs757343	Iranian	260 (55), 181 (57)	PCR-RFLP	N	(55, 57)		

(Continued)

**Table 1** Continued.

Gene	Description	Polymorphism	Population	Number of PCOS patients	Methods	Association	Ref.
<i>MTHFR</i>	Methylenetetrahydrofolate reductase	C677T	Italian, Korean, Turkish, Iranian, Indian, Polish	70 (73), 227 (75), 86 (76), 38 (77), 92 (78), 98 (79), 76 (80), 261 (81)	PCR-RFLP, real-time PCR	N	(73, 75, 76, 77, 78, 79, 80, 81)
		A1298C	Iranian, Polish	38, 76	PCR-RFLP, None	Y, N	(77), (80)
<i>PPAR-γ</i>	Peroxisome proliferator-activated receptor gamma	Pro12Ala	European, Egyptian	1598, 100	Meta-analysis, PCR-RFLP	N, Y	(87), (88)
<i>IL-1β/IL-1Ra</i>	Interleukin-1beta/IL-1 receptor antagonist	IL-1β (C[-511] T)	Indian	95	PCR-RFLP	N	(95)
		IL-1Ra (VNTR within intron 2)	Indian	95		N	(95)

GWAS, genome-wide association study; N, gene polymorphisms show none significant association with PCOS; PCR-RFLP, polymerase chain reaction-restriction fragment-length polymorphism assay; TADA, TaqMan allelic discrimination assay; Y, gene polymorphisms show significant association with PCOS.

Participation of *MTHFR* polymorphisms in the regulation of the folate pathway or decreased activity of the homocysteine pathway has been revealed in a variety of pathologies, including Alzheimer's disease, neural tube defects, acute leukemia, colon cancer and PCOS (67). Laboratory data indicate that variate C677T in the *MTHFR* gene decreases the activity of the folate pathway by 50% (59, 68, 69). However, the few clinical studies done to date have obtained similar results that indicate the association between *MTHFR* and PCOS (70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81). More recently, Wang *et al.* (82) conducted a meta-analysis to evaluate the effect of *MTHFR* C677T polymorphism on the risk of PCOS. The results from the analysis of 14 articles involving 1478 PCOS cases from 16 distinct populations showed that the *MTHFR* C667T variant had different roles in the risks of PCOS in different ethnicities. The T allele significantly increased the risk of PCOS in Asians, had a protective effect in Caucasians and had no significant associations in Middle Eastern individuals.

### PPAR-γ polymorphism

PCOS is related to some extent with hormonal and metabolic disorders, with insulin resistance being one of the most common metabolic disturbances (83). The *PPAR-γ* (*PPARG*) gene participates in the management of insulin sensitivity and in lipid metabolism (84). The gene product (*PPAR-γ*) plays an important part in the management of energy storage and insulin sensitivity based on its roles in glucose regulating and lipid metabolism (85). Wang *et al.* (86) investigated

the association of *PPARG* and *PPARG-1alpha* gene polymorphisms with PCOS among 201 Chinese Han PCOS women and 147 controls. The results suggested that these gene polymorphisms were not associated with the risk of PCOS. A meta-analysis conducted in 2010 evaluated the relationship of Pro12Ala polymorphism of *PPARG* with PCOS (87). The results from the analysis of 13 studies that included 1598 PCOS cases and 1881 controls revealed no significant association of the Pro12Ala polymorphism with PCOS. However, in a more recent investigation of the association of *PPARG* Pro12Ala polymorphism with PCOS, the frequency of the polymorphic allele was significantly higher in PCOS cases than in the controls, suggesting that the *PPARG* Pro12Ala polymorphism might contribute to the risk of PCOS (88).

### IL-1β/IL-1Ra gene polymorphisms

PCOS is recognized as pro-inflammatory condition (89). Elevation of circulating inflammatory mediators have been observed in PCOS (90), including pro-inflammatory cytokines, white blood cells, chemokines and multiple endothelial inflammation markers (91). Cytokines are important in inflammatory processes and are also thought to be associated with the etiology of PCOS (92). IL-1β is a member of the IL-1 family; it is a valid pro-inflammatory cytokine that has a great impact on the physiology of reproduction. IL-1β has also been reported as a crucial moderating element in ovulation, fertilization, embryo implantation and tissue restructuring (93, 94). Another important member of the IL-1 family is IL-1RA; it may

be able to minimize the inflammatory response. Various *IL-1Ra* gene polymorphisms have been identified and six distinct alleles have been recognized to date. A recent study by Rashid *et al.* (95) involving 95 women with PCOS and 45 healthy women as controls analyzed the connection between *IL-1β* and *IL-1Ra* gene polymorphisms with the risk of PCOS. The findings indicated that polymorphisms in these cytokines genes may contribute to the risk of PCOS.

## Conclusions

PCOS is a common endocrine disorder among women of reproductive aged. Many risk factors affect the occurrence and development of PCOS, including ovary abnormalities, obesity and environmental factors. Multiple studies have suggested that genetic factors are essential in the etiology of PCOS. As we have described, a number of polymorphic variants of candidate genes associated with PCOS have been identified in case-control association studies (Table 1). However, the genetic significance and the exact pathophysiological relevance of most of these genes with PCOS, which include *StAR*, *FSHR* and *FTO*, are still unclear and need to be confirmed. For the putative candidate genes identified so far, more studies are needed for a better understanding of their molecular and physiological roles in PCOS. These data will clarify the risk factors of PCOS and contribute to the prediction of the disorder, which could improve diagnosis and treatment of PCOS patients. Polymorphic variants of genes have been garnered mainly by genome-wide analysis and epigenetic regulation such as DNA methylation and noncoding RNAs have been more recently linked with PCOS. The advances in epigenetic studies on PCOS have been reviewed elsewhere (96).

The etiology of PCOS remains difficult to clarify because of heterogeneities in its features and origins. With the systematic acquisition of more data, identification of better phenotypes and studies involving larger populations, the pathological mechanisms of PCOS will become better understood, which will further improve the diagnosis and knowledge of the clinical manifestations of PCOS. These data will also help us determine whether one genetic polymorphism has a compelling association with PCOS. Several candidate genes have been associated with the PCOS susceptibility. However, controversial findings still exist, which may ascribe to the diverse populations and genotyping methods. There is a long way to go before we entirely understand the genetic effects on this complex and heterogeneous disorder. Further studies

should consider address genetic and environment factors, including obesity and nutrition, which will provide a better understanding of the etiology of PCOS.

### Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

### Funding

This work was supported by the program for the development of county key medical disciplines of Zhejiang Province and the key medical disciplines funds in Hangzhou.

### Author contribution statement

Yao Chen was the primary author of this manuscript. She reviewed and edited the manuscript and approved the final version. Shu-ying Fang contributed to the writing and formatting of this work.

## References

- Nandi A, Chen Z, Patel R & Poretsky L. Polycystic ovary syndrome. *Endocrinology Metabolism Clinics of North America* 2014 **43** 123–147. (<https://doi.org/10.1016/j.ecl.2013.10.003>)
- Wang Y, Zhao X, Zhao H, Ding H, Tan J, Chen J, Zhang R, Azziz R & Yang D. Risks for gestational diabetes mellitus and pregnancy-induced hypertension are increased in polycystic ovary syndrome. *BioMed Research International* 2013 **2013** 182582. (<https://doi.org/10.1155/2013/182582>)
- Sirmans SM & Pate KA. Epidemiology, diagnosis, and management of polycystic ovary syndrome. *Clinical Epidemiology* 2013 **6** 1–13. (<https://doi.org/10.2147/CLEP.S37559>)
- Hardiman P, Pillay OC & Atiomo W. Polycystic ovary syndrome and endometrial carcinoma. *Lancet* 2003 **361** 1810–1812. (<https://doi.org/S0140673603134095>)
- Segars JH & DeCherney AH. Is there a genetic basis for polycystic ovary syndrome? *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 2058–2060. (<https://doi.org/10.1210/jc.2010-0518>)
- Vink JM, Sadrzadeh S, Lambalk CB & Boomsma DI. Heritability of polycystic ovary syndrome in a Dutch twin-family study. *Journal of Clinical Endocrinology and Metabolism* 2006 **91** 2100–2104. (<https://doi.org/10.1210/jc.2005-1494>)
- Hague WM, Adams J, Reeders ST, Peto TE & Jacobs HS. Familial polycystic ovaries: a genetic disease? *Clinical Endocrinology* 1988 **29** 593–605. (<https://doi.org/10.1111/j.1365-2265.1988.tb03707.x>)
- Lunde O, Magnus P, Sandvik L & Hoglo S. Familial clustering in the polycystic ovarian syndrome. *Gynecologic and Obstetric Investigation* 1989 **28** 23–30. (<https://doi.org/10.1159/000293493>)
- Kahsar-Miller MD, Nixon C, Boots LR, Go RC & Azziz R. Prevalence of polycystic ovary syndrome (PCOS) in first-degree relatives of patients with PCOS. *Fertility and Sterility* 2001 **75** 53–58. ([https://doi.org/S0015-0282\(00\)01662-9](https://doi.org/S0015-0282(00)01662-9) [pii])
- Legro RS, Driscoll D, Strauss JF 3rd, Fox J & Dunaif A. Evidence for a genetic basis for hyperandrogenemia in polycystic ovary syndrome. *PNAS* 1998 **95** 14956–14960. (<https://doi.org/10.1073/pnas.95.25.14956>)
- Colilla S, Cox NJ & Ehrmann DA. Heritability of insulin secretion and insulin action in women with polycystic ovary syndrome and their first degree relatives. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 2027–2031. (<https://doi.org/10.1210/jcem.86.5.7518>)

- 12 Menke MN & Strauss JF 3rd. Genetic approaches to polycystic ovarian syndrome. *Current Opinion in Obstetrics and Gynecology* 2007 **19** 355–359. (<https://doi.org/10.1097/GCO.0b013e328220e877>)
- 13 Mykhalchenko K, Lizneva D, Trofimova T, Walker W, Suturina L, Diamond MP & Azziz R. Genetics of polycystic ovary syndrome. *Expert Review of Molecular Diagnostics* 2017 **17** 723–733. (<https://doi.org/10.1080/14737159.2017.1340833>)
- 14 Barbar E, Lehoux JG & Lavigne P. Toward the NMR structure of StAR. *Molecular and Cellular Endocrinology* 2009 **300** 89–93. (<https://doi.org/10.1016/j.mce.2008.12.007>)
- 15 Miller WL. Steroidogenic acute regulatory protein (StAR), a novel mitochondrial cholesterol transporter. *Biochimica et Biophysica Acta* 2007 **1771** 663–676. (<https://doi.org/10.1016/j.bbali.2007.02.012>)
- 16 Kahsar-Miller MD, Conway-Myers BA, Boots LR & Azziz R. Steroidogenic acute regulatory protein (StAR) in the ovaries of healthy women and those with polycystic ovary syndrome. *American Journal of Obstetrics and Gynecology* 2001 **185** 1381–1387. (<https://doi.org/10.1067/mob.2001.118656>)
- 17 Jakubowski L. Genetic aspects of polycystic ovary syndrome. *Endokrynologia Polska* 2005 **56** 285–293.
- 18 Nazouri AS, Khosravifar M, Akhlaghi AA, Shiva M & Afsharian P. No relationship between most polymorphisms of steroidogenic acute regulatory (StAR) gene with polycystic ovarian syndrome. *International Journal of Reproductive BioMedicine* 2015 **13** 771–778. (<https://doi.org/10.29252/ijrm.13.12.771>)
- 19 Kosova G & Urbanek M. Genetics of the polycystic ovary syndrome. *Molecular and Cellular Endocrinology* 2013 **373** 29–38. (<https://doi.org/10.1016/j.mce.2012.10.009>)
- 20 Naor Z. Signaling by G-protein-coupled receptor (GPCR): studies on the GnRH receptor. *Frontiers in Neuroendocrinology* 2009 **30** 10–29. (<https://doi.org/10.1016/j.yfrne.2008.07.001>)
- 21 Torrealday S, Lalioti MD, Guzeloglu-Kayisli O & Seli E. Characterization of the gonadotropin releasing hormone receptor (GnRHR) expression and activity in the female mouse ovary. *Endocrinology* 2013 **154** 3877–3887. (<https://doi.org/10.1210/en.2013-1341>)
- 22 Aguilar-Rojas A & Huerta-Reyes M. Human gonadotropin-releasing hormone receptor-activated cellular functions and signaling pathways in extra-pituitary tissues and cancer cells (Review). *Oncological Reports* 2009 **22** 981–990.
- 23 Chen WY, Du YQ, Guan X, Zhang HY & Liu T. Effect of GnRHR polymorphisms on in vitro fertilization and embryo transfer in patients with polycystic ovary syndrome. *Journal of Human Genetics* 2017 **62** 1065–1071. (<https://doi.org/10.1038/jhg.2017.85>)
- 24 Caburet S, Fruchter RB, Legois B, Fellous M, Shalev S & Veitia RA. A homozygous mutation of GnRHR in a familial case diagnosed with polycystic ovary syndrome. *European Journal of Endocrinology* 2017 **176** K9–K14. (<https://doi.org/10.1530/EJE-16-0968>)
- 25 Kim JJ, Choi YM, Hong MA, Chae SJ, Hwang K, Yoon SH, Ku SY, Suh CS & Kim SH. FSH receptor gene p. Thr307Ala and p. Asn680Ser polymorphisms are associated with the risk of polycystic ovary syndrome. *Journal of Assisted Reproduction and Genetics* 2017 **34** 1087–1093. (<https://doi.org/10.1007/s10815-017-0953-z>)
- 26 Layman LC, Amde S, Cohen DP, Jin M & Xie J. The Finnish follicle-stimulating hormone receptor gene mutation is rare in North American women with 46,XX ovarian failure. *Fertility and Sterility* 1998 **69** 300–302. ([https://doi.org/10.1016/S0015-0282\(97\)00480-9](https://doi.org/10.1016/S0015-0282(97)00480-9))
- 27 Simoni M, Gromoll J & Nieschlag E. The follicle-stimulating hormone receptor: biochemistry, molecular biology, physiology, and pathophysiology. *Endocrine Reviews* 1997 **18** 739–773. (<https://doi.org/10.1210/edrv.18.6.0320>)
- 28 Perez Mayorga M, Gromoll J, Behre HM, Gassner C, Nieschlag E & Simoni M. Ovarian response to follicle-stimulating hormone (FSH) stimulation depends on the FSH receptor genotype. *Journal of Clinical Endocrinology and Metabolism* 2000 **85** 3365–3369. (<https://doi.org/10.1210/jcem.85.9.6789>)
- 29 Laven JS, Mulders AG, Suryandari DA, Gromoll J, Nieschlag E, Fauser BC & Simoni M. Follicle-stimulating hormone receptor polymorphisms in women with normogonadotropic anovulatory infertility. *Fertility and Sterility* 2003 **80** 986–992. ([https://doi.org/10.1016/S0015-0282\(03\)01115-4](https://doi.org/10.1016/S0015-0282(03)01115-4))
- 30 Greb RR, Grieshaber K, Gromoll J, Sonntag B, Nieschlag E, Kiesel L & Simoni M. A common single nucleotide polymorphism in exon 10 of the human follicle stimulating hormone receptor is a major determinant of length and hormonal dynamics of the menstrual cycle. *Journal of Clinical Endocrinology and Metabolism* 2005 **90** 4866–4872. (<https://doi.org/10.1210/jc.2004-2268>)
- 31 de Koning CH, Benjamins T, Harms P, Homburg R, van Montfrans JM, Gromoll J, Simoni M & Lambalk CB. The distribution of FSH receptor isoforms is related to basal FSH levels in subfertile women with normal menstrual cycles. *Human Reproduction* 2006 **21** 443–446. (<https://doi.org/10.1093/humrep/dei317>)
- 32 Overbeek A, Kuijper EA, Hendriks ML, Blankenstein MA, Ketel IJ, Twisk JW, Hompes PG, Homburg R & Lambalk CB. Clomiphene citrate resistance in relation to follicle-stimulating hormone receptor Ser680Ser-polymorphism in polycystic ovary syndrome. *Human Reproduction* 2009 **24** 2007–2013. (<https://doi.org/10.1093/humrep/dep114>)
- 33 Valkenburg O, Uitterlinden AG, Piersma D, Hofman A, Themmen AP, de Jong FH, Fauser BC & Laven JS. Genetic polymorphisms of GnRH and gonadotrophic hormone receptors affect the phenotype of polycystic ovary syndrome. *Human Reproduction* 2009 **24** 2014–2022. (<https://doi.org/10.1093/humrep/dep113>)
- 34 Fu L, Zhang Z, Zhang A, Xu J, Huang X, Zheng Q, Cao Y, Wang L & Du J. Association study between FSHR Ala307Thr and Ser680Asn variants and polycystic ovary syndrome (PCOS) in Northern Chinese Han women. *Journal of Assisted Reproduction and Genetics* 2013 **30** 717–721. (<https://doi.org/10.1007/s10815-013-9979-z>)
- 35 Hubacek JA, Stanek V, Gebauerova M, Pilipincova A, Dlouha D, Poledne R, Aschermann M, Skalicka H, Matouskova J, Kruger A, et al. A FTO variant and risk of acute coronary syndrome. *Clinica Chimica Acta* 2010 **411** 1069–1072. (<https://doi.org/10.1016/j.cca.2010.03.037>)
- 36 Frayling TM, Timpson NJ, Weedon MN, Zeggini E, Freathy RM, Lindgren CM, Perry JR, Elliott KS, Lango H, Rayner NW, et al. A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* 2007 **316** 889–894. (<https://doi.org/10.1126/science.1141634>)
- 37 Willer CJ, Speliotes EK, Loos RJ, Li S, Lindgren CM, Heid IM, Berndt SI, Elliott AL, Jackson AU, Lamina C, et al. Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. *Nature Genetics* 2009 **41** 25–34. (<https://doi.org/10.1038/ng.287>)
- 38 Demirci H, Yilmaz M, Ergun MA, Yurtcu E, Bukan N & Ayvaz G. Frequency of adiponectin gene polymorphisms in polycystic ovary syndrome and the association with serum adiponectin, androgen levels, insulin resistance and clinical parameters. *Gynecological Endocrinology* 2010 **26** 348–355. (<https://doi.org/10.3109/09513590903367051>)
- 39 Shi Y, Guo M, Yan J, Sun W, Zhang X, Geng L, Xu L & Chen Z. Analysis of clinical characteristics in large-scale Chinese women with polycystic ovary syndrome. *Neuro Endocrinology Letters* 2007 **28** 807–810.
- 40 Yan Q, Hong J, Gu W, Zhang Y, Liu Q, Su Y, Zhang Y, Li X, Cui B & Ning G. Association of the common rs9939609 variant of FTO gene with polycystic ovary syndrome in Chinese women. *Endocrine* 2009 **36** 377–382. (<https://doi.org/10.1007/s12020-009-9257-0>)
- 41 Barber TM, Bennett AJ, Groves CJ, Sovio U, Ruokonen A, Martikainen H, Pouta A, Hartikainen AL, Elliott P, Lindgren CM, et al. Association of variants in the fat mass and obesity associated (FTO) gene with polycystic ovary syndrome. *Diabetologia* 2008 **51** 1153–1158. (<https://doi.org/10.1007/s00125-008-1028-6>)

- 42 Li T, Wu K, You L, Xing X, Wang P, Cui L, Liu H, Cui Y, Bian Y, Ning Y, *et al.* Common variant rs9939609 in gene FTO confers risk to polycystic ovary syndrome. *PLoS ONE* 2013 **8** e66250. (<https://doi.org/10.1371/journal.pone.0066250>)
- 43 Ramos RB & Spritzer PM. FTO gene variants are not associated with polycystic ovary syndrome in women from Southern Brazil. *Gene* 2015 **560** 25–29. (<https://doi.org/10.1016/j.gene.2015.01.012>)
- 44 Kim JJ, Choi YM, Hong MA, Kim JM, Hwang SS, Lee GH, Chae SJ, Hwang KR, Yoon SH & Kim SH. Gene dose effect between a fat mass and obesity-associated polymorphism and body mass index was observed in Korean women with polycystic ovary syndrome but not in control women. *Fertility and Sterility* 2014 **102** 1143.e1142–1148. e1142. (<https://doi.org/10.1016/j.fertnstert.2014.07.004>)
- 45 Saxena R & Welt CK. Polycystic ovary syndrome is not associated with genetic variants that mark risk of type 2 diabetes. *Acta Diabetologica* 2013 **50** 451–457. (<https://doi.org/10.1007/s00592-012-0383-4>)
- 46 Ewens KG, Jones MR, Ankener W, Stewart DR, Urbanek M, Dunaif A, Legro RS, Chua A, Azziz R, Spielman RS, *et al.* FTO and MC4R gene variants are associated with obesity in polycystic ovary syndrome. *PLoS ONE* 2011 **6** e16390. (<https://doi.org/10.1371/journal.pone.0016390>)
- 47 Cai X, Liu C & Mou S. Association between fat mass- and obesity-associated (FTO) gene polymorphism and polycystic ovary syndrome: a meta-analysis. *PLoS ONE* 2014 **9** e86972. (<https://doi.org/10.1371/journal.pone.0086972>)
- 48 Liu AL, Xie HJ, Xie HY, Liu J, Yin J, Hu JS & Peng CY. Association between fat mass and obesity associated (FTO) gene rs9939609 A/T polymorphism and polycystic ovary syndrome: a systematic review and meta-analysis. *BMC Medical Genetics* 2017 **18** 89. (<https://doi.org/10.1186/s12881-017-0452-1>)
- 49 Bagheri M, Abdi-Rad I, Hosseini-Jazani N, Zarrin R, Nanbakhsh F & Mohammadzaie N. An association study between INSR/NsiI (rs2059806) and INSR/PmlI (rs1799817) SNPs in women with polycystic ovary syndrome from West Azerbaijan Province, Iran. *Journal of Reproduction and Infertility* 2015 **16** 109–112.
- 50 Shi X, Xie X, Jia Y & Li S. Associations of insulin receptor and insulin receptor substrates genetic polymorphisms with polycystic ovary syndrome: a systematic review and meta-analysis. *Journal of Obstetrics and Gynaecology Research* 2016 **42** 844–854. (<https://doi.org/10.1111/jog.13002>)
- 51 Reis GV, Gontijo NA, Rodrigues KE, Alves MT, Ferreira CN & Gomes KB. Vitamin D receptor polymorphisms and the polycystic ovary syndrome: a systematic review. *Journal of Obstetrics and Gynaecology Research* 2017 **43** 436–446. (<https://doi.org/10.1111/jog.13250>)
- 52 Haussler MR, Jurutka PW, Mizwicki M & Norman AW. Vitamin D receptor (VDR)-mediated actions of 1 $\alpha$ ,25(OH) $_2$ vitamin D(3): genomic and non-genomic mechanisms. *Best Practice and Research Clinical Endocrinology and Metabolism* 2011 **25** 543–559. (<https://doi.org/10.1016/j.beem.2011.05.010>)
- 53 Ogunkolade BW, Boucher BJ, Prahil JM, Bustin SA, Burrin JM, Noonan K, North BV, Mannan N, McDermott ME, DeLuca HF, *et al.* Vitamin D receptor (VDR) mRNA and VDR protein levels in relation to vitamin D status, insulin secretory capacity, and VDR genotype in Bangladeshi Asians. *Diabetes* 2002 **51** 2294–2300. (<https://doi.org/10.2337/diabetes.51.7.2294>)
- 54 Mahmoudi T. Genetic variation in the vitamin D receptor and polycystic ovary syndrome risk. *Fertility and Sterility* 2009 **92** 1381–1383. (<https://doi.org/10.1016/j.fertnstert.2009.05.002>)
- 55 Zadeh-Vakili A, Ramezani Tehrani F, Daneshpour MS, Zarkesh M, Saadat N & Azizi F. Genetic polymorphism of vitamin D receptor gene affects the phenotype of PCOS. *Gene* 2013 **515** 193–196. (<https://doi.org/10.1016/j.gene.2012.11.049>)
- 56 El-Shal AS, Shalaby SM, Aly NM, Rashad NM & Abdelaziz AM. Genetic variation in the vitamin D receptor gene and vitamin D serum levels in Egyptian women with polycystic ovary syndrome. *Molecular Biology Reports* 2013 **40** 6063–6073. (<https://doi.org/10.1007/s11033-013-2716-y>)
- 57 Ranjanz F, Mahmoudi T, Irani Shemirani A, Mahban A, Nikzamir A, Vahedi M, Ashrafi M & Gourabi H. A common variant in the adiponectin gene and polycystic ovary syndrome risk. *Molecular Biology Reports* 2012 **39** 2313–2319. (<https://doi.org/10.1007/s11033-011-0981-1>)
- 58 Jedrzejuk D, Laczanski L, Milewicz A, Kuliczowska-Plaksej J, Lenarcik-Kabza A, Hirnle L, Zaleska-Dorobisz U & Lwow F. Classic PCOS phenotype is not associated with deficiency of endogenous vitamin D and VDR gene polymorphisms rs731236 (TaqI), rs7975232 (ApaI), rs1544410 (BsmI), rs10735810 (FokI): a case-control study of lower Silesian women. *Gynecological Endocrinology* 2015 **31** 976–979. (<https://doi.org/10.3109/09513590.2015.1062865>)
- 59 Yamada K, Chen Z, Rozen R & Matthews RG. Effects of common polymorphisms on the properties of recombinant human methylenetetrahydrofolate reductase. *PNAS* 2001 **98** 14853–14858. (<https://doi.org/10.1073/pnas.261469998>)
- 60 Reilly R, McNulty H, Pentieva K, Strain JJ & Ward M. MTHFR 677TT genotype and disease risk: is there a modulating role for B-vitamins? *Proceedings of the Nutrition Society* 2014 **73** 47–56. (<https://doi.org/10.1017/S0029665113003613>)
- 61 Frosst P, Blom HJ, Milos R, Goyette P, Sheppard CA, Matthews RG, Boers GJ, den Heijer M, Kluijtmans LA, van den Heuvel LP, *et al.* A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nature Genetics* 1995 **10** 111–113. (<https://doi.org/10.1038/ng0595-111>)
- 62 Willmott M, Bartosik DB & Romanoff EB. The effect of folic acid on superovulation in the immature rat. *Journal of Endocrinology* 1968 **41** 439–445. (<https://doi.org/10.1677/joe.0.0410439>)
- 63 Mohanty D & Das KC. Effect of folate deficiency on the reproductive organs of female rhesus monkeys: a cytomorphological and cytokinetic study. *Journal of Nutrition* 1982 **112** 1565–1576. (<https://doi.org/10.1093/jn/112.8.1565>)
- 64 Badawy A, State O, El Gawad S & El Aziz OA. Plasma homocysteine and polycystic ovary syndrome: the missed link. *European Journal of Obstetrics and Gynecology and Reproductive Biology* 2007 **131** 68–72. (<https://doi.org/10.1016/j.ejogrb.2006.10.015>)
- 65 Kazerooni T, Asadi N, Dehbashi S & Zolghadri J. Effect of folic acid in women with and without insulin resistance who have hyperhomocysteinemic polycystic ovary syndrome. *International Journal of Gynecology and Obstetrics* 2008 **101** 156–160. (<https://doi.org/10.1016/j.ijgo.2007.10.024>)
- 66 Bayraktar F, Dereli D, Ozgen AG & Yilmaz C. Plasma homocysteine levels in polycystic ovary syndrome and congenital adrenal hyperplasia. *Endocrine Journal* 2004 **51** 601–608. (<https://doi.org/10.1507/endocrj.51.601>)
- 67 Trimmer EE. Methylenetetrahydrofolate reductase: biochemical characterization and medical significance. *Current Pharmaceutical Design* 2013 **19** 2574–2593. (<https://doi.org/10.2174/1381612811319140008>)
- 68 Cortese C & Motti C. MTHFR gene polymorphism, homocysteine and cardiovascular disease. *Public Health Nutrition* 2001 **4** 493–497.
- 69 McQuillan BM, Beilby JP, Nidorf M, Thompson PL & Hung J. Hyperhomocysteinemia but not the C677T mutation of methylenetetrahydrofolate reductase is an independent risk determinant of carotid wall thickening. The Perth Carotid Ultrasound Disease Assessment Study (CUDAS). *Circulation* 1999 **99** 2383–2388. (<https://doi.org/10.1161/01.CIR.99.18.2383>)
- 70 Glueck CJ, Wang P, Fontaine RN, Sieve-Smith L, Tracy T & Moore SK. Plasminogen activator inhibitor activity: an independent risk factor for the high miscarriage rate during pregnancy in women with polycystic ovary syndrome. *Metabolism* 1999 **48** 1589–1595. ([https://doi.org/10.1016/S0026-0495\(99\)90250-0](https://doi.org/10.1016/S0026-0495(99)90250-0))
- 71 Sills ES, Genton MG, Perloe M, Schattman GL, Bralley JA & Tucker MJ. Plasma homocysteine, fasting insulin, and androgen patterns among women with polycystic ovaries and infertility. *Journal of Obstetrics and Gynaecology Research* 2001 **27** 163–168. (<https://doi.org/10.1111/j.1447-0756.2001.tb01241.x>)



- 72 Tsanadis G, Vartholomatos G, Korkontzelos I, Avgoustatos F, Kakosimos G, Sotiriadis A, Tatsioni A, Eleftheriou A & Lolis D. Polycystic ovarian syndrome and thrombophilia. *Human Reproduction* 2002 **17** 314–319. (<https://doi.org/10.1093/humrep/17.2.314>)
- 73 Orio F Jr, Palomba S, Di Biase S, Colao A, Tauchmanova L, Savastano S, Labella D, Russo T, Zullo F & Lombardi G. Homocysteine levels and C677T polymorphism of methylenetetrahydrofolate reductase in women with polycystic ovary syndrome. *Journal of Clinical Endocrinology and Metabolism* 2003 **88** 673–679. (<https://doi.org/10.1210/jc.2002-021142>)
- 74 Palep-Singh M, Picton HM, Yates ZR, Barth J & Balen AH. Polycystic ovary syndrome and the single nucleotide polymorphisms of methylenetetrahydrofolate reductase: a pilot observational study. *Human Fertility* 2007 **10** 33–41. (<https://doi.org/10.1080/14647270600950157>)
- 75 Choi SW, Gu BH, Ramakrishna S, Park JM & Baek KH. Association between a single nucleotide polymorphism in MTHFR gene and polycystic ovary syndrome. *European Journal of Obstetrics and Gynecology and Reproductive Biology* 2009 **145** 85–88. (<https://doi.org/10.1016/j.ejogrb.2009.04.013>)
- 76 Karadeniz M, Erdogan M, Zengi A, Eroglu Z, Tamsel S, Olukman M, Saygili F & Yilmaz C. Methylenetetrahydrofolate reductase C677T gene polymorphism in Turkish patients with polycystic ovary syndrome. *Endocrine* 2010 **38** 127–133. (<https://doi.org/10.1007/s12020-010-9370-0>)
- 77 Idali F, Zareii S, Mohammad-Zadeh A, Reihany-Sabet F, Akbarzadeh-Pasha Z, Khorram-Khorshid HR, Zarnani AH & Jeddi-Tehrani M. Plasminogen activator inhibitor 1 and methylenetetrahydrofolate reductase gene mutations in Iranian women with polycystic ovary syndrome. *American Journal of Reproductive Immunology* 2012 **68** 400–407. (<https://doi.org/10.1111/aji.12002>)
- 78 Jain M, Pandey P, Tiwary NK & Jain S. MTHFR C677T polymorphism is associated with hyperlipidemia in women with polycystic ovary syndrome. *Journal of Human Reproductive Sciences* 2012 **5** 52–56. (<https://doi.org/10.4103/0974-1208.97802>)
- 79 Ozegowska K, Bogacz A, Bartkowiak-Wieczorek J, Seremak-Mrozikiewicz A & Pawelczyk L. Is there an association between the development of metabolic syndrome in PCOS patients and the C677T MTHFR gene polymorphism? *Ginekologia Polska* 2016 **87** 246–253. (<https://doi.org/10.17772/gp/61751>)
- 80 Szafarowska M, Segiet A & Jerzak MM. Methylenetetrahydrofolate reductase A1298C and C677T polymorphisms and adverse pregnancy outcome in women with PCOS. *Neuro Endocrinology Letters* 2016 **37** 141–146.
- 81 Carlus SJ, Sarkar S, Bansal SK, Singh V, Singh K, Jha RK, Sadasivam N, Sadasivam SR, Giresha PS, Thangaraj K, et al. Is MTHFR 677 C>T polymorphism clinically important in polycystic ovarian syndrome (PCOS)? A case-control study, meta-analysis and trial sequential analysis. *PLoS ONE* 2016 **11** e0151510. (<https://doi.org/10.1371/journal.pone.0151510>)
- 82 Wang L, Xu W, Wang C, Tang M & Zhou Y. Methylenetetrahydrofolate reductase C677T polymorphism and the risks of polycystic ovary syndrome: an updated meta-analysis of 14 studies. *Oncotarget* 2017 **8** 59509–59517. (<https://doi.org/10.18632/oncotarget.18472>)
- 83 Farrell K & Antoni MH. Insulin resistance, obesity, inflammation, and depression in polycystic ovary syndrome: biobehavioral mechanisms and interventions. *Fertility and Sterility* 2010 **94** 1565–1574. (<https://doi.org/10.1016/j.fertnstert.2010.03.081>)
- 84 Falcao-Pires I, Goncalves N, Henriques-Coelho T, Moreira-Goncalves D, Roncon-Albuquerque R Jr & Leite-Moreira AF. Apelin decreases myocardial injury and improves right ventricular function in monocrotaline-induced pulmonary hypertension. *American Journal of Physiology: Heart and Circulatory Physiology* 2009 **296** H2007–H2014. (<https://doi.org/10.1152/ajpheart.00089.2009>)
- 85 Auwerx J. PPARgamma, the ultimate thrifty gene. *Diabetologia* 1999 **42** 1033–1049. (<https://doi.org/10.1007/s001250051268>)
- 86 Wang S, Fu C, Wang H, Shi Y, Xu X, Chen J, Song X, Sun K, Wang J, Fan X, et al. Polymorphisms of the peroxisome proliferator-activated receptor-gamma coactivator-1alpha gene are associated with hypertrophic cardiomyopathy and not with hypertension hypertrophy. *Clinical Chemistry and Laboratory Medicine* 2007 **45** 962–967. (<https://doi.org/10.1515/CCLM.2007.189>)
- 87 Tang ST, Wang CJ, Tang HQ, Peng WJ, Wang YM & Zhang Q. Association of Pro12Ala polymorphism in peroxisome proliferator-activated receptor gamma with polycystic ovary syndrome: a meta-analysis. *Molecular Biology Reports* 2012 **39** 9649–9660. (<https://doi.org/10.1007/s11033-012-1830-6>)
- 88 Zaki M, Hassan N, El-Bassyouni HT, Kamal S, Basha W, Azmy O & Amr K. Association of the Pro12Ala polymorphism with the metabolic parameters in women with polycystic ovary syndrome. *Open Access Macedonian Journal of Medical Sciences* 2017 **5** 275–280. (<https://doi.org/10.3889/oamjms.2017.088>)
- 89 Shorakae S, Teede H, de Courten B, Lambert G, Boyle J & Moran LJ. The emerging role of chronic low-grade inflammation in the pathophysiology of polycystic ovary syndrome. *Seminars in Reproductive Medicine* 2015 **33** 257–269. (<https://doi.org/10.1055/s-0035-1556568>)
- 90 Jatzko B & Ott J. Circulating inflammatory markers in polycystic ovary syndrome: a systematic review and meta-analysis. *Fertility and Sterility* 2011 **96** e158. (<https://doi.org/10.1016/j.fertnstert.2011.07.1098>)
- 91 Papalou O, Livadas S, Karachalios A, Tolia N, Kokkoris P, Tripolitakis K & Diamanti-Kandarakis E. White blood cells levels and PCOS: direct and indirect relationship with obesity and insulin resistance, but not with hyperandrogenemia. *Hormones* 2015 **14** 91–100. (<https://doi.org/10.14310/horm.2002.1563>)
- 92 Ebejer K & Calleja-Agius J. The role of cytokines in polycystic ovarian syndrome. *Gynecological Endocrinology* 2013 **29** 536–540. (<https://doi.org/10.3109/09513590.2012.760195>)
- 93 Bonetti TC, Salomao R, Brunialti M, Braga DP, Borges E Jr & Silva ID. Cytokine and hormonal profile in serum samples of patients undergoing controlled ovarian stimulation: interleukin-1beta predicts ongoing pregnancy. *Human Reproduction* 2010 **25** 2101–2106. (<https://doi.org/10.1093/humrep/deq171>)
- 94 Prutsch N, Fock V, Haslinger P, Haider S, Fiala C, Pollheimer J & Knofler M. The role of interleukin-1beta in human trophoblast motility. *Placenta* 2012 **33** 696–703. (<https://doi.org/10.1016/j.placenta.2012.05.008>)
- 95 Rashid N, Nigam A, Saxena P, Jain SK & Wajid S. Association of IL-1beta, IL-1Ra and FABP1 gene polymorphisms with the metabolic features of polycystic ovary syndrome. *Inflammation Research* 2017 **66** 621–636. (<https://doi.org/10.1007/s00011-017-1045-3>)
- 96 Li S, Zhu D, Duan H & Tan Q. The epigenomics of polycystic ovarian syndrome: from pathogenesis to clinical manifestations. *Gynecological Endocrinology* 2016 **32** 942–946. (<https://doi.org/10.1080/09513590.2016.1203409>)

Received in final form 27 March 2018

Accepted 5 April 2018

Accepted Preprint published online 5 April 2018

