

REVIEW

The impact of endocrine disruptor chemicals on oocyte/embryo and clinical outcomes in IVF

Rosalie Cabry¹, Philippe Merviel², Aicha Madkour^{1,3}, Elodie Lefranc¹, Florence Scheffler¹, Rachel Desailoud¹, Véronique Bach¹ and Moncef Benkhalifa¹

¹Amiens University, Amiens, Haut-de-France, France

²Brest University, Brest, Bretagne, France

³Mohammed V University of Rabat, Reproductive Medicine, Rabat, Morocco

Correspondence should be addressed to M Benkhalifa: benkhalifamoncef78@gmail.com

Abstract

The negative impact of endocrine-disrupting pesticides on human fertility is now a key issue in reproductive health. There are much fewer literature data about the impact of pesticide exposure on women than on men and very few studies of women participating in an *in vitro* fertilization (IVF) programme. In the present review, we found that (1) various pesticides with an endocrine-disrupting action are associated with poor oocyte maturation and competency, embryonic defects and poor IVF outcomes, and (2) some pesticide compounds are linked to specific causes of female infertility, such as premature ovarian insufficiency, polycystic ovarian syndrome, and endometriosis. IVF participants living in agricultural regions should be informed about the fertility decline, low ongoing pregnancy rates, and elevated risk of miscarriage associated with exposure to high doses of pesticides.

Key Words

- ▶ endocrine disruptor
- ▶ oocyte-embryo
- ▶ clinical outcome
- ▶ IVF

Endocrine Connections
(2020) **9**, R134–R142

Introduction

Infertility is defined as failure to obtain a clinical pregnancy after 12 months of regular, unprotected sexual intercourse. On average, it affects 8–12% of couples of child-bearing age (1). A decline in human fertility has prompted an increasing proportion of couples to enrol in *in vitro* fertilization (IVF) programmes. Over the last 50 years, the sperm count has fallen by 32–50% in Europe and United States (2, 3); this fall is too rapid to be due to a genetic factor but might be related to one or more environmental factors, such as exposure to pesticides.

Pesticide are substances or combinations of substances used in many areas of agriculture and industry. They include (1) phytosanitary products (also referred to as phytopharmaceuticals) used in agricultural or non-agricultural plant sectors to control insects (insecticides), weeds (herbicides), fungi and moulds (fungicides), or pests (rodenticides); (2) biocides used in industry (treatment of wood and textiles), hospital environments (hydroalcoholic gels), and domestic settings (disinfection); and (3)

antiparasitics used in human and veterinary medicine (treatment of lice, scabies, mites, fleas, ticks, etc.) (4, 5).

The main families of pesticides by chemical composition are as follows:

- **Organophosphorus** (OP) compounds, such as chlorpyrifos, cypermethrin, malathion, and parathion
- **Organochlorine** (OC) compounds, such as methoxychlor (MXC), 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), dichlorodiphenyldichloroethane (DDD), dichlorodiphenyldichloroethylene (DDE), dichlorodiphenyltrichloroethane (DDT), polychlorinated biphenyls (PCBs), dieldrin, endosulfan, hexachlorobenzene (HCB), hexachlorocyclohexane (HCH), and lindane
- **Carbamates**, such as carbofuran, benomyl, and mancozeb
- **Pyrethroids**, such as permethrin
- **Triazines**, such as atrazine

Many pesticides are known to be endocrine-disrupting chemicals (EDCs), defined as 'exogenous agents, that are potentially capable of synthesis, secretion, transport, binding, action, or elimination of the natural hormones responsible for the maintenance of homeostasis, reproduction, and developmental processes in the body' (5). Even though the involvement of EDCs in certain reproductive diseases is well documented, only few studies have examined the direct impact of pesticides on fertility (and especially in female infertility) in IVF programmes. Hence, the objective of the present review was to assess the impact of the most frequently used pesticides on female fertility, oocyte-embryo quality, and clinical outcomes in IVF programmes.

Pesticides as EDCs

A large body of evidence from animal studies and epidemiological surveys shows that pesticides like bisphenol A (BPA) and phthalates have an endocrine-disrupting action and a reprotoxic impact (5, 6, 7, 8, 9, 10, 11). Furthermore, Manikkam (12) highlighted the existence of a transgenerational risk in the rat: *in utero* exposure of pups to a mixture of pesticides (administered by gavaging the pregnant dams) was associated with changes in puberty onset, spermatogenesis, and the development of ovarian follicles up to the F3 generation.

The direct, toxic endocrine-disrupting effect (i.e. a dose-response relationship) has rarely been characterized for a single target compound. In general, the major toxic exposure is associated with a large number of low-dose compounds and the interactions between these compounds. This 'cocktail effect' is why the presence of even very small amounts of some specific pesticides can perturb an organism's hormonal balance (13, 14). Indeed, some pesticides bind as agonists to hormone receptors, which results in direct cell damage and/or the dysregulation of one or more biochemical or genetic pathways. In turn, this dysregulation produces toxic metabolites and increases the level of oxidative stress (14), which affects both male and female fertility. There are more research data on the influence of various pesticides on male fertility than on female fertility. However, even the data on male fertility (with putative links between exposure and a range of disorders), especially those correlated with poor sperm quality and low testicular weight (15, 16, 17), make it hard to affirm the presence of a direct, causal effect in humans.

Compounds such as lindane, PCB, atrazine, and mancozeb might dysregulate hormonal status in women by decreasing LH concentrations and thus promoting oligo-ovulation/anovulation and follicle destruction (13, 14).

Pesticide exposure and female fertility

In 1994, De Cock *et al.* (18) investigated the impact of pesticide exposure in market gardeners in the Netherlands and found that the fecundability ratio fell as the intensity of pesticide exposure increased. In this study, 28% of couples in the 'high exposure' group and 8% of the couples in 'low exposure' group requested assisted reproductive technology (ART) support. Other studies have emphasized this effect of pesticide exposure in women, with lengthening of the time required to conceive (19, 20, 21, 22). A risk of spontaneous miscarriage has also been observed, particularly in Bretveld *et al.*'s (23) study of female farmers (odds ratio (95% CI)=4.0 (1.14–14.01).

Data from the Agricultural Health Study cohort showed that menstrual cycles of women exposed to pesticides increased in length, although no distinction was made between occupational exposure and domestic exposure (24).

These results clearly converge on an impact of pesticide use (especially in an occupational setting) on female reproductive capacity and certain female infertility diseases, such as premature ovarian failure (POF), polycystic ovarian syndrome (PCOS), and endometriosis. Furthermore, pesticide exposure might have direct or indirect effects on oocyte/embryo quality and the clinical outcomes of IVF.

The association between pesticide exposure and certain female infertility diseases

The negative impact of pesticides has been investigated in animal experiments. Many deficiencies have been described, such as low ovarian weight (25, 26), impaired folliculogenesis (27), a high aneuploidy rate, and the acceleration of follicular atresia (25, 26, 27).

Indeed, women exposed to some endocrine-disrupting pesticides (such as atrazine, lindane, and maneb) have an elevated risk of long menstrual cycles or anovulation (13). Conversely, high blood levels of DDE (28) and DDT (29) are associated with luteal phase deficiency and short menstrual cycles. The most serious consequence is POF (Premature Ovarian failure), defined as the cessation

of menstrual cycles before the age of 40 years and characterized by very low blood levels of anti-Müllerian hormone (AMH). The prevalence of POF is around 1%, as a result of genetic or autoimmune/metabolic factors or cancer therapy (1). Nonetheless, POF might sometimes be caused by environmental factors, including pesticide exposure. Indeed, a few studies have linked exposure to HCH, mirex (30), pyrethroids (31), and their metabolites (especially 3-phenoxybenzoic acid (32)), PCB, and DDT (33) to ovarian ageing and menopause at an earlier average age. Interestingly, low AMH levels were associated with the presence of DDT in the serum (detected in more than 40% of samples from patients with POF (33)) and with high concentrations of pyrethroids and their metabolites with 25–26% reduction (31, 32).

The presence of DDE, MXC, or simazine (a triazine herbicide) was associated with follicular atresia (34). Given that MXC, simazine, and other pesticides (as atrazine, endosulfan, and chlordecone) are oestrogen receptor agonists (in contrast to DDE, DDT, and vinclozolin), they may have oestrogenic and/or antiandrogenic effects (5, 10, 35, 36). This implicitly explains why some specific pesticides could be responsible for various female reproductive diseases to differing extents and why the effects of some pesticide compounds (such as DDE, PCB, and MXC) depend on the patient's genetic predisposition. Indeed, these pesticides were associated with abnormally high AMH levels and a high number of small antral follicles – reflecting the presence of PCOS (37, 38, 39). Polycystic ovarian syndrome is a heterogeneous condition that affects 5–10% of women. According to the Rotterdam diagnostic criteria for PCOS, two of the following three criteria must be met: infrequent or absent ovulation (oligospomenorrhea), an abnormal ovarian morphology on ultrasound, and hyperandrogenism (1).

Moreover, some pesticides (such as TCDD (40, 41, 42), PCB (43), and permethrin (44)) have a documented impact on the endometrium by increasing vascularization, cell proliferation, VEGF levels, and thus endometriosis (with prevalence of 0.8–6%) (1). However, other pesticides (cypermethrin (45), chlorpyrifos, malathion, diazinon, lindane, DDT, and pyrethroids (46)) have the opposite effect by indirectly inhibiting endometrial proliferation or causing oxidative damage to the uterus.

Many studies have found an association between pesticide concentrations in the follicular fluid on one hand and the number and quality of oocytes collected for IVF (47, 48), endometrial thickness, and embryo implantation rates (46) on the other. Studies of animal models have shown that pesticide exposure is associated

with elevated oxidative stress, which may be responsible for the observed alterations (49).

Impact on oocyte quality

Exposure to pesticides might cause generalized oxidative stress (49), with the elevated production of free radicals and impacts on superoxide dismutase, catalase, glutathione peroxidase, glutathione reductase, and glutathione transferase. This might explain why oocyte quality is impacted; oocyte dysmorphisms may reflect poor competence for development into a good embryo for implantation.

Indeed, we have highlighted (50) a conclusive link between a high prevalence (>75%) of oocytes with centrally located cytoplasm granulation (CLCG) and the patients' exposure to pesticides in the highly agricultural Picardy area of northern France (annual pesticide consumption: ~3900 tons). As a result, ongoing pregnancy and live birth rates were lower in couples with a high prevalence of oocytes with CLCG than in couples with low (<25%) prevalence (14% vs 32% for the ongoing pregnancy rate and 13% vs 30% for the live birth rate, respectively). Conversely, the early miscarriage rate was higher (47%) in the high-prevalence group than in the low-prevalence group (11%; odds ratio: 3.1). This poor IVF outcome might be indirectly due to high levels of pesticide exposure (over 3000 g/ha), which engenders a higher risk of oocytes with CLCG.

In line with these results, some pesticides (e.g. atrazine, cypermethrin, DDT, dieldrin, MCX, and vinclozolin) affect folliculogenesis (via the granulosa and theca cells) and thus induce meiotic aberrations (aneuploidy) and follicular atresia (51). For example, DDT stimulates aromatase and acts in synergy with FSH to induce a premature rise in oestradiol levels, affecting oocyte maturation (51). Furthermore, PCB might damage the ovarian reserve, the follicular response to administered gonadotropins, and oocyte maturation (51). Indeed, exposure of immature porcine cumulus-oocyte complexes to an organochlorine mixture (48) or atrazine (52) during *in vitro* maturation was associated with an elevated incidence of incompletely matured oocytes. The researchers mentioned the following pathophysiological mechanisms: abnormal cellular shape (disrupted spindle morphology), abnormal mitochondrial activity, and alterations in oocyte DNA (48, 51, 52).

The pesticides most frequently linked to defective oocyte maturation and oocyte competence are malathion (53), parathion (54), MXC (55), DDD/DDE/DDT/PCB

Table 1 Summary of the negative impact of various pesticides on oocytes, embryos, the endometrium, and clinical outcomes in IVF.

Pesticide	EReg	Oocyte quality	Oocyte maturation	Oocyte genetic	Embryo development	Endometrium	IR	CPR	References
Organophosphorus									
Chlorpyrifos			+		+	+	+		(71)
Cypermethrin			+			+			(46)
Malathion	+		+		+	+			(45)
Parathion	+	+	+	+					(53)
									(46)
									(80)
									(10)
									(54)
Organochlorine									
MXC	+	+	+	+					(27)
									(55)
TCDD					++	++	+	+	(81)
									(49)
									(40)
									(41)
									(42)
DDD	++								(56)
DDE	++	+	+		++	+			(28)
									(56)
									(39)
DDT	++	+	+	++	+	+	+	+	(26)
									(29)
									(56)
									(31)
									(20)
									(47)
									(51)
PCB	++	+			++	++	+++	++	(43)
									(56)
									(77)
									(78)
									(39)
									(72)
Diieldrin	+				++				(58)
									(71)
Endosulfan						++	+++		(79)
									(57)
HCB	++						++	+	(47)
HCH	++	+	+					+	(30)
									(47)
Lindane	+	++	++	+		+	+		(13)
									(59)
									(46)

Carbamate	Carbofuran	+	+++	+	+	(82)
	Benomyl	++	+	++		(61) (73)
Pyrethroid	Mancozeb	+	+	+	++	(62) (60) (44)
	Permethrin	+				(51) (13)
Triazine	Atrazine	++	+	+	+	(36) (52)
Vinclozoline		+++				(83) (51)

The '+' symbols indicate a negative impact: '+' rarely reported, with a low level of interest; '++': widely reported, with a high level of interest. CPR: clinical pregnancy rate; DDD: dichlorodiphenyldichloroethane; DDE: dichlorodiphenyldichloroethylene; DDT: dichlorodiphenyltrichloroethane; EReg: endocrine regulation; HCB: hexachlorobenzene; HCH: hexachlorocyclohexane; IR: implantation rate; MXC: methoxychlor; PCB: polychlorinated biphenyl; TCDD: 2,3,7,8-tetrachlorodibenzo-p-dioxin.

(56, 57), dieldrin (58), HCB/HCH (47), lindane (59), benomyl (60), and mancozeb (61, 62) (Table 1).

Interestingly, DDT was associated with a low proportion of diploid oocytes (56), while benomyl interfered with microtubules (60) and lindane was associated with cell damage (especially vacuolation and cytoplasmic fragmentation) (59). It was reported that mancozeb can strongly affect the meiotic spindle organization of oocytes (61), but the degree of damage could be decreased by treatment with resveratrol (62).

Moreover, exposure to deltamethrin (another common insecticide) in a mouse model was linked to cellular oxidative stress and meiotic abnormalities via DNA damage (63). Liu *et al.* (49) suggested that MXC exposure induces oxidative stress and affects mouse oocyte meiotic maturation via the accumulation of superoxide radicals and other reactive oxygen species (ROS), aberrant mitochondrial distribution, a low mitochondrial membrane potential, and elevated lipid peroxidation. Thus, exposure to MXC can negatively affect oocyte meiotic maturation – primarily through impairments in cellular metabolism. In general, all these pesticide-linked defects in oocyte quality are likely to degrade competence for the development of a genetically undamaged embryo.

Impact on embryo development

Pesticides can affect indirectly the embryo by dysregulating embryonic genome activation and embryonic metabolism, which is dependent on oxygen uptake. This latter is low at the 8-cell stage (64, 65, 66) and tends to increase after the morula stage required for blastocyst expansion (67). Blastocyst formation and the number of cells per blastocyst declined with the concentration of organochlorines (48) and atrazine (52). Furthermore, the association of pesticides like deltamethrin (63) and MXC (49) with poor metabolic and genetic status during embryo development might be due to the impairment of various biochemical pathways (13) and high ROS production. Moreover, high ROS concentrations and generalized oxidative stress are likely to affect the integrity of cellular constituents, such as DNA and proteins (68, 69, 70).

Various pesticides (especially chlorpyrifos with endosulfan (71), malathion (53), DDE/DDT/PCB (56, 47), dieldrin (72), mancozeb (73) (Table 1), and pretilachlor/diazinon (46)) may impact negatively embryonic development. Consequently, the use of low-quality embryos associated with pesticide exposure would give poor clinical outcomes (e.g. an elevated risk of miscarriage) in IVF programmes.

Impact on clinical outcomes in IVF programmes

Women who consume high levels of pesticide residues in fruits and vegetables (74) or who live in an area with high pesticide exposure (50, 75) have an above-average risk of miscarriage. In one study, the probability of clinical pregnancy was 18% below average and the live birth rate was 26% below average in the women most exposed to pesticides (74). However, in a study in California, there were no differences in terms of spontaneous miscarriage, preeclampsia, and preterm birth rates between women exposed to pesticides (occupationally or through residence in an agricultural area) and unexposed women (76). Pesticides associated with poor clinical outcomes in IVF are chlorpyrifos (46), TCDD (42), DDT/PCB (46, 56, 77, 78), HCB/HCH (47, 56, 57), and endosulfan (79) (Table 1). Lindane, DDT, diazinon, and chlorpyrifos were associated with a low implantation rate but did not have a clear impact on the clinical pregnancy and live birth rates (46). Nevertheless, PCB and endosulfan have been linked to repeated implantation failures (77, 79).

Conclusion

A growing number of studies have assessed the putative causal link between exposure to pesticides and female fertility disorders. Although awareness of these issues has increased, the literature data are too scarce for conclusive, decisive recommendations. The impact of exposure to various endocrine-disrupting pesticides on fertility is now a public health issue that urgently requires the performance of more epidemiological studies – especially those focused on female fertility and women in IVF programmes. Furthermore, it is important to design studies that assess the severity of exposure and the nature of pesticides. There is also a need to develop more specific, rapid diagnosis techniques and treatments that might decrease pesticide-induced damage. Indeed, IVF participants living in agricultural regions should be informed about the fertility decline, low ongoing pregnancy rates, and elevated risk of miscarriage associated with exposure to high doses of pesticides.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was funded by Amiens-Picardie University Hospital (Amiens, France) and Jules Verne University of Picardie (Peritox Laboratory, Amiens, France).

Acknowledgements

The authors thank the staff at Amiens-Picardie University Hospital's ART unit and the staff at the Peritox Laboratory for their support.

References

- Vander Borgh M & Wyns C. Fertility and infertility: definition and epidemiology. *Clinical Biochemistry* 2018 **62** 2–10. (<https://doi.org/10.1016/j.clinbiochem.2018.03.012>)
- Manikkam M, Tracey R, Guerrero-Bosagna C & Skinner MK. Pesticide and insect repellent mixture (permethrin and DEET) induces epigenetic transgenerational inheritance of disease and sperm epimutations. *Reproductive Toxicology* 2012 **34** 708–719. (<https://doi.org/10.1016/j.reprotox.2012.08.010>)
- Levine H, Jørgensen N, Martino-Andrade A, Mendiola J, Weksler-Derri D, Mindlis I, Pinotti R & Swan SH. Temporal trends in sperm count: a systematic review and meta-regression analysis. *Human Reproduction Update* 2017 **23** 646–659. (<https://doi.org/10.1093/humupd/dmx022>)
- Sengupta P, Borges E, Dutta S & Krajewska-Kulak E. Decline in sperm count in European men during the past 50 years. *Human and Experimental Toxicology* 2018 **37** 247–255. (<https://doi.org/10.1177/0960327117703690>)
- Alavanja MC. Introduction: pesticides use and exposure, extensive worldwide. *Reviews on Environmental Health* 2009 **24** 303–309. (<https://doi.org/10.1515/reveh.2009.24.4.303>)
- Sifakis S, Mparmpas M, Soldin OP & Tsatsakis A. Pesticide exposure and health related issues in male and female reproductive system. In *Pesticides-Formulations, Effects, Fate*. London, UK: Intech Open, 2011. (<https://doi.org/10.5772/15845>)
- Rocheleau CM, Romitti PA & Dennis LK. Pesticides and hypospadias: a meta-analysis. *Journal of Pediatric Urology* 2009 **5** 17–24. (<https://doi.org/10.1016/j.jpuro.2008.08.006>)
- Carmichael SL, Yang W, Roberts EM, Kegley SE, Wolff C, Guo L, Lammer EJ, English P & Shaw GM. Hypospadias and residential proximity to pesticide applications. *Pediatrics* 2013 **132** e1216–e1226. (<https://doi.org/10.1542/peds.2013-1429>)
- Haraux E, Tourneux P, Kouakam C, Stephan-Blanchard E, Boudailliez B, Leke A, Klein C & Chardon K. Isolated hypospadias: the impact of prenatal exposure to pesticides, as determined by meconium analysis. *Environment International* 2018 **119** 20–25. (<https://doi.org/10.1016/j.envint.2018.06.002>)
- Mnif W, Pillon A, Balaguer P & Bartegi A. Endocrine xenoestrogens disrupters, molecular mechanisms and detection methods. *Therapie* 2007 **62** 369–386. (<https://doi.org/10.2515/therapie:2007062>)
- Bhardwaj JK & Saraf P. Granulosa cell apoptosis by impairing antioxidant defense system and cellular integrity in caprine antral follicles post Malathion exposure. *Environmental Toxicology* 2016 **31** 1944–1954. (<https://doi.org/10.1002/tox.22195>)
- Ma Y, He X, Qi K, Wang T, Qi Y, Cui L, Wang F & Song M. Effects of environmental contaminants on fertility and reproductive health. *Journal of Environmental Sciences* 2019 **77** 210–217. (<https://doi.org/10.1016/j.jes.2018.07.015>)
- Farr SL, Cooper GS, Cai J, Savitz DA & Sandler DP. Pesticide use and menstrual cycle characteristics among premenopausal women in the Agricultural Health Study. *American Journal of Epidemiology* 2004 **160** 1194–1204. (<https://doi.org/10.1093/aje/kwi006>)
- Bretveld RW, Thomas CM, Scheepers PT, Zielhuis GA & Roelveland N. Pesticide exposure: the hormonal function of the female reproductive system disrupted? *Reproductive Biology and Endocrinology* 2006 **4** 30. (<https://doi.org/10.1186/1477-7827-4-30>)
- Joshi SC, Mathur R & Gulati N. Testicular toxicity of chlorpyrifos (an organophosphate pesticide) in albino rat. *Toxicology and Industrial Health* 2007 **23** 439–444. (<https://doi.org/10.1177/0748233707080908>)

- 16 Akhtar N, Srivastava MK & Raizada RB. Assessment of chlorpyrifos toxicity on certain organs in rat, *Rattus norvegicus*. *Journal of Environmental Biology* 2009 **30** 1047–1053.
- 17 Farag AT, Radwan AH, Sorour F, El Okazy A, El-Agamy el-S & El-Sebae Ael-K. Chlorpyrifos induced reproductive toxicity in male mice. *Reproductive Toxicology* 2010 **29** 80–85. (<https://doi.org/10.1016/j.reprotox.2009.10.003>)
- 18 De Cock J, Westveer K, Heederik D, te Velde E & van Kooij R. Time to pregnancy and occupational exposure to pesticides in fruit growers in the Netherlands. *Occupational and Environmental Medicine* 1994 **51** 693699. (<https://doi.org/10.1136/oem.51.10.693>)
- 19 Curtis KM, Savitz DA, Weinberg CR & Arbuckle TE. The effect of pesticide exposure on time to pregnancy. *Epidemiology* 1999 **10** 112117. (<https://doi.org/10.1097/00001648-199903000-00005>)
- 20 Harley KG, Marks AR, Bradman A, Barr DB & Eskenazi B. DDT exposure, work in agriculture, and time to pregnancy among farmworkers in California. *Journal of Occupational and Environmental Medicine* 2008 **50** 13351342. (<https://doi.org/10.1097/JOM.0b013e31818f684d>)
- 21 Idrovo AJ, Sanin LH, Cole D, Chavarro J, Cáceres H, Narváez J & Restrepo M. Time to first pregnancy among women working in agricultural production. *International Archives of Occupational and Environmental Health* 2005 **78** 493500. (<https://doi.org/10.1007/s00420-005-0615-9>)
- 22 Abell A, Juul S & Bonde JPE. Time to pregnancy among female greenhouse workers. *Scandinavian Journal of Work, Environment and Health* 2000 **26** 131136. (<https://doi.org/10.5271/sjweh.522>)
- 23 Bretveld RW, Hooiveld M, Zielhuis GA, Pellegrino A, van Rooij IALM & Roeleveld N. Reproductive disorders among male and female greenhouse workers. *Reproductive Toxicology* 2008 **25** 107–114. (<https://doi.org/10.1016/j.reprotox.2007.08.005>)
- 24 Alavanja MC, Sandler DP, McMaster SB, Zahm SH, McDonnell CJ, Lynch CF, Pennybacker M, Rothman N, Dosemeci M, Bond AE, *et al.* The agricultural health study. *Environmental Health Perspectives* 1996 **104** 362369. (<https://doi.org/10.1289/ehp.96104362>)
- 25 Park S, Kim S, Jin H, Lee K & Bae J. Impaired development of female mouse offspring maternally exposed to simazine. *Environmental Toxicology and Pharmacology* 2014 **38** 845–851. (<https://doi.org/10.1016/j.etap.2014.10.008>)
- 26 El-Sharkawy EE, Kames AO, Sayed SM, Nisr NA, Wahba NM, Elsherif WM, Nafady AM, Abdel-Hafeez MM & Aamer AA. The ameliorative effect of propolis against methoxychlor induced ovarian toxicity in rat. *Experimental and Toxicologic Pathology* 2014 **66** 415–421. (<https://doi.org/10.1016/j.etp.2014.06.003>)
- 27 Satar DA, Tap O & Ay MO. Electron microscopic examination of the effects of methyl parathion exposure on the ovaries. *European Review for Medical and Pharmacological Sciences* 2015 **19** 2725–2731.
- 28 Windham GC, Lee D, Mitchell P, Anderson M, Petreas M & Lasley B. Exposure to organochlorine compounds and effects on ovarian function. *Epidemiology* 2005 **16** 182–190. (<https://doi.org/10.1097/01.ede.0000152527.24339.17>)
- 29 Ouyang F, Perry MJ, Venners SA, Chen C, Wang B, Yang F, Fang Z, Zang T, Wang L, Xu X, *et al.* Serum DDT, age at menarche, and abnormal menstrual cycle length. *Occupational and Environmental Medicine* 2005 **62** 878–884. (<https://doi.org/10.1136/oem.2005.020248>)
- 30 Grindler NM, Allsworth JE, Macones GA, Kannan K, Roehl KA & Cooper AR. Persistent organic pollutants and early menopause in U.S. Women. *PLoS ONE* 2015 **10** e0116057. (<https://doi.org/10.1371/journal.pone.0116057>)
- 31 Whitworth KW, Baird DD, Steiner AZ, Bornman RM, Travlos GS, Wilson RE & Longnecker MP. AntiMüllerian hormone and lifestyle, reproductive, and environmental factors among women in rural South Africa. *Epidemiology* 2015 **26** 429–435. (<https://doi.org/10.1097/EDE.0000000000000265>)
- 32 Li C, Cao M, Ma L, Ye X, Song Y, Pan W, Xu Z, Ma X, Lan Y, Chen P, *et al.* Pyrethroid pesticide exposure and risk of primary ovarian insufficiency in Chinese women. *Environmental Science and Technology* 2018 **52** 3240–3248. (<https://doi.org/10.1021/acs.est.7b06689>)
- 33 Pan W, Ye X, Yin S, Ma X, Li C, Zhou J, Liu W & Liu J. Selected persistent organic pollutants associated with the risk of primary ovarian insufficiency in women. *Environment International* 2019 **129** 51–58. (<https://doi.org/10.1016/j.envint.2019.05.023>)
- 34 Vabre P, Gatimel N, Moreau J, Gayrard V, Picard-Hagen N, Parinaud J & Leandri RD. Environmental pollutants, a possible etiology for premature ovarian insufficiency: a narrative review of animal and human data. *Environmental Health* 2017 **16** 37. (<https://doi.org/10.1186/s12940-017-0242-4>)
- 35 Kadhel P, Monnier P, Boucoiran I, Chaillet N & Fraser WD. Organochlorine pollutants and female fertility: a systematic review focusing on in vitro fertilization studies. *Reproductive Sciences* 2012 **19** 1246–1259. (<https://doi.org/10.1177/1933719112446077>)
- 36 Samardzija D, Pogrmic-Majkic K, Fa S, Glisic B, Stanic B & Andric N. Atrazine blocks ovulation via suppression of Lhr and Cyp19a1 mRNA and estradiol secretion in immature gonadotropin-treated rats. *Reproductive Toxicology* 2016 **61** 10–18. (<https://doi.org/10.1016/j.reprotox.2016.02.009>)
- 37 Manikkam M, Haque MM, Guerrero-Bosagna C, Nilsson EE & Skinner MK. Pesticide methoxychlor promotes the epigenetic transgenerational inheritance of adult-onset disease through the female germline. *PLoS ONE* 2014 **9** e102091. (<https://doi.org/10.1371/journal.pone.0102091>)
- 38 Zhang J, Liu XF, Liu Y, Xu LZ, Zhou LL, Tang LL, Zhuang J, Li TT, Guo WQ, Hu R, *et al.* Environmental risk factors for women with polycystic ovary syndrome in china: a population-based case-control study. *Journal of Biological Regulators and Homeostatic Agents* 2014 **28** 203–211.
- 39 Yang Q, Zhao Y, Qiu X, Zhang C, Li R & Qiao J. Association of serum levels of typical organic pollutants with polycystic ovary syndrome (PCOS): a case-control study. *Human Reproduction* 2015 **30** 1964–1973. (<https://doi.org/10.1093/humrep/dev123>)
- 40 Bofinger DP, Feng L, Chi LH, Love J, Stephen FD, Sutter TR, Osteen KG, Costich TG, Batt RE, Koury ST, *et al.* Effect of TCDD exposure on CYP1A1 and CYP1B1 expression in explant cultures of human endometrium. *Toxicological Sciences* 2001 **62** 299–314. (<https://doi.org/10.1093/toxsci/62.2.299>)
- 41 Nayyar T, Bruner-Tran KL, Piestrzeniewicz-Ulanska D & Osteen KG. Developmental exposure of mice to TCDD elicits a similar uterine phenotype in adult animals as observed in women with endometriosis. *Reproductive Toxicology* 2007 **23** 326–336. (<https://doi.org/10.1016/j.reprotox.2006.09.007>)
- 42 Foster WG. Endocrine toxicants including 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and dioxin-like chemicals and endometriosis: is there a link? *Journal of Toxicology and Environmental Health* 2008 **11** 177–187.
- 43 Jirsova S, Masata JV, Drbohlava P, Pavelkova J, Jech L, Omelka M & Zvárová J. Differences in the polychlorinated biphenyl levels in follicular fluid in individual types of sterility. *Ceska Gynekologie* 2005 **70** 262–268.
- 44 Noori E, Nasri S, Janan A, Mohebbi A, Moini A, Ramazani F & Aflatoonian R. Expression of vascular endothelial growth factor receptors in endometriosis. *International Journal of Fertility and Sterility* 2013 **7** 114.
- 45 Al-Hamdani NMH & Yajurvedi HN. Effect of cypermethrin on the ovarian activity and its impact on fertility and pubertal onset of offspring. *Beni-Suef University Journal of Basic and Applied Sciences* 2017 **6** 374–382. (<https://doi.org/10.1016/j.bjbas.2017.07.003>)
- 46 Al Hussaini TK, Addelaleem AA, Elnashar I, Shabaan OM, Mostafa R, El-Baz MAH, El-Deek SEM & Farghaly TA. The effect of follicular fluid pesticides and polychlorinated biphenyls concentrations on intracytoplasmic sperm injection (ICSI) embryological and

- clinical outcome. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* 2018 **220** 39–43. (<https://doi.org/10.1016/j.ejogrb.2017.11.003>)
- 47 Petro EM, Leroy JL, Covaci A, Franssen E, De Neubourg D, Dirtu AC, De Pauw I & Bols PE. Endocrine-disrupting chemicals in human follicular fluid impair in vitro oocyte developmental competence. *Human Reproduction* 2012 **27** 1025–1033. (<https://doi.org/10.1093/humrep/der448>)
- 48 Campagna C, Sirad MA, Ayotte P & Bailey JL. Impaired maturation, fertilization and embryonic development of porcine oocytes following exposure to an environmentally relevant organochlorine mixture. *Biology of Reproduction* 2001 **65** 554–560. (<https://doi.org/10.1095/biolreprod65.2.554>)
- 49 Liu Y, Wang YL, Chen MH, Zhang Z, Xu BH, Liu R, Xu L, He SW, Li FP, Qi ZQ, *et al.* Methoxychlor exposure induces oxidative stress and affects mouse oocyte meiotic maturation. *Molecular Reproduction and Development* 2016 **83** 768–779. (<https://doi.org/10.1002/mrd.22683>)
- 50 Merviel P, Cabry R, Chardon K, Haraux E, Scheffler F, Mansouri NB, Devaux A, Chahine H, Bach V, Copin H, *et al.* Impact of oocytes with CLCG on ICSI outcomes and their potential relation to pesticide exposure. *Journal of Ovarian Research* 2017 **10** 42. (<https://doi.org/10.1186/s13048-017-0335-2>)
- 51 Marques-Pinto A & Carvalho D. Human infertility: are endocrine disruptors to blame? *Endocrine Connections* 2013 **2** R15–R29. (<https://doi.org/10.1530/EC-13-0036>)
- 52 Yuan B, Liang S, Jin Y-X, Zhang M-J, Zhang JB & Kim N-H. Toxic effects of atrazine on porcine oocytes and possible mechanisms of action. *PLoS ONE* 2017 **22** e0179861. (<https://doi.org/10.1371/journal.pone.0179861>)
- 53 Zhou YJ, Huang HR, Zhou J & Wang LQ. beta-Cypermethrin exposure affects female reproduction by enhancing oxidative stress in mice uterine tissue. *Regulatory Toxicology and Pharmacology* 2018 **98** 284–290. (<https://doi.org/10.1016/j.yrtph.2018.08.011>)
- 54 Aziz F, Mehboob Z, Jalali S, Shami SA & Qureshi IZ. Effect of organophosphate pesticide Malathion on in-vitro oocyte maturation, fertilization and embryo development of nili ravi buffalo (*Bubalus bubalis*). In *Proceedings of 6th Asian Buffalo Congress (Vol. 27, p. 30)*, 2009.
- 55 Armenti AE, Zama AM, Passantino L & Uzumcu M. Developmental methoxychlor exposure affects multiple reproductive parameters and ovarian folliculogenesis and gene expression in adult rats. *Toxicology and Applied Pharmacology* 2008 **233** 286–296. (<https://doi.org/10.1016/j.taap.2008.09.010>)
- 56 Jirsová S, Mašata J, Jech L & Zvárová J. Effect of polychlorinated biphenyls (PCBs) and 1,1,1-trichloro-2,2-bis (4-chlorophenyl)-ethane (DDT) in follicular fluid on the results of in vitro fertilization-embryo transfer (IVF-ET) programs. *Fertility and Sterility* 2010 **93** 1831–1836. (<https://doi.org/10.1016/j.fertnstert.2008.12.063>)
- 57 Mahalingaiah S, Missmer SA, Maity A, Williams PL, Meeker JD, Berry K, Ehrlich S, Perry MJ, Cramer DW & Hauser R. Association of hexachlorobenzene (HCB), dichlorodiphenyltrichloroethane (DDT), and dichlorodiphenyldichloroethylene (DDE) with in vitro fertilization (IVF) outcomes. *Environmental Health Perspectives* 2012 **120** 316–320. (<https://doi.org/10.1289/ehp.1103696>)
- 58 Bastos AMX, Souza Mdo C, Almeida Filho GLD, Krauss TM, Pavesi T & Silva LED. Organochlorine compound levels in fertile and infertile women from Rio de Janeiro, Brazil. *Arquivos Brasileiros de Endocrinologia e Metabologia* 2013 **57** 346–353. (<https://doi.org/10.1590/s0004-27302013000500003>)
- 59 Palmerini MG, Zhurabekova G, Balmagambetova A, Nottola SA, Miglietta S, Belli M, Bianchi S, Cecconi S, Di Nisio V, Familiari G, *et al.* The pesticide lindane induces dose-dependent damage to granulosa cells in an in vitro culture. *Reproductive Biology* 2017 **17** 349–356. (<https://doi.org/10.1016/j.repbio.2017.09.008>)
- 60 Morinaga H, Yanase T, Nomura M, Okabe T, Goto K, Harada N & Nawata H. A benzimidazole fungicide, benomyl, and its metabolite, carbendazim, induce aromatase activity in a human ovarian granulosa-like tumor cell line (KGN). *Endocrinology* 2004 **145** 1860–1869. (<https://doi.org/10.1210/en.2003-1182>)
- 61 Rossi G, Palmerini MG, Macchiarelli G, Buccione R & Cecconi S. Mancozeb adversely affects meiotic spindle organization and fertilization in mouse oocytes. *Reproductive Toxicology* 2006 **22** 51–55. (<https://doi.org/10.1016/j.reprotox.2005.11.005>)
- 62 Liu Y, Wang YL, He SW, Chen MH, Zhang Z, Fu XP, Fu BB, Liao BQ, Lin YH, Qi ZQ, *et al.* Protective effects of resveratrol against mancozeb induced apoptosis damage in mouse oocytes. *Oncotarget* 2017 **8** 6233–6245. (<https://doi.org/10.18632/oncotarget.14056>)
- 63 Hia ZZ, Zhang JW, Zhou D, Xu DQ & Feng XZ. Deltamethrin exposure induces oxidative stress and affects meiotic maturation in mouse oocyte. *Chemosphere* 2019 **223** 704–713. (<https://doi.org/10.1016/j.chemosphere.2019.02.092>)
- 64 Houghton FD, Thompson JG, Kennedy CJ & Leese HJ. Oxygen consumption and energy metabolism of the early mouse embryo. *Molecular Reproduction and Development* 1996 **44** 476–485. ([https://doi.org/10.1002/\(SICI\)1098-2795\(199608\)44:4<476::AID-MRD7>3.0.CO;2-I](https://doi.org/10.1002/(SICI)1098-2795(199608)44:4<476::AID-MRD7>3.0.CO;2-I))
- 65 Trimarchi JR, Liu L, Porterfield DM, Smith PJ & Keefe DL. Oxidative phosphorylation-dependent and independent oxygen consumption by individual preimplantation mouse embryos. *Biology of Reproduction* 2000 **62** 1866–1874. (<https://doi.org/10.1095/biolreprod62.6.1866>)
- 66 Thompson JG, Partridge RJ, Houghton FD, Cox CI & Leese HJ. Oxygen uptake and carbohydrate metabolism by in vitro derived bovine embryos. *Journal of Reproduction and Fertility* 1996 **106** 299–306. (<https://doi.org/10.1530/jrf.0.1060299>)
- 67 Donnay I & Leese HJ. Embryo metabolism during the expansion of the bovine blastocyst. *Molecular Reproduction and Development* 1999 **53** 171–178. ([https://doi.org/10.1002/\(SICI\)1098-2795\(199906\)53:2<171::AID-MRD6>3.0.CO;2-F](https://doi.org/10.1002/(SICI)1098-2795(199906)53:2<171::AID-MRD6>3.0.CO;2-F))
- 68 Leese HJ. Metabolism of the preimplantation embryo: 40 years on. *Reproduction* 2012 **143** 417–427. (<https://doi.org/10.1530/REP-11-0484>)
- 69 Guerin P, El Mouatassim S & Menezes Y. Oxidative stress and protection against reactive oxygen species in the pre-implantation embryo and its surroundings. *Human Reproduction Update* 2001 **7** 175–189. (<https://doi.org/10.1093/humupd/7.2.175>)
- 70 Nasr-Esfahani MH, Aitken JR & Johnson MH. Hydrogen peroxide levels in mouse oocytes and early cleavage stage embryos developed in vitro or in vivo. *Development* 1990 **109** 501–507.
- 71 Nandi S, Gupta PSP, Roy SC, Selvaraju S & Ravindra JP. Chlorpyrifos and endosulfan affect buffalo oocyte maturation, fertilization, and embryo development in vitro directly and through cumulus cells. *Environmental Toxicology* 2011 **26** 57–67. (<https://doi.org/10.1002/tox.20529>)
- 72 Berger T & Horner CM. In vivo exposure of female rats to toxicants may affect oocyte quality. *Reproductive Toxicology* 2003 **17** 273–281. ([https://doi.org/10.1016/s0890-6238\(03\)00009-1](https://doi.org/10.1016/s0890-6238(03)00009-1))
- 73 Cecconi S, Paro R, Rossi G & Macchiarelli G. The effects of the endocrine disruptors dithiocarbamates on the mammalian ovary with particular regard to mancozeb. *Current Pharmaceutical Design* 2007 **13** 2989–3004. (<https://doi.org/10.2174/138161207782110516>)
- 74 Chiu YH, Williams PL, Gillman MW, Gaskins AJ, Mingez-Alarcon L, Souter I, Toth TL, Ford JB, Hauser R, Chavarro JE, *et al.* Association between pesticide residue intake from consumption of fruits and vegetables and pregnancy outcomes among women undergoing infertility treatment with assisted reproductive technology. *JAMA Internal Medicine* 2018 **178** 17–26. (<https://doi.org/10.1001/jamainternmed.2017.5038>)
- 75 Arbuckle TE, Lin Z & Mery LS. An exploratory analysis of the effect of pesticide exposure on the risk of spontaneous abortion in an

- Ontario farm population. *Environmental Health Perspectives* 2001 **109** 851–857. (<https://doi.org/10.1289/ehp.01109851>)
- 76 Willis WO, de Peyster A, Molgaard CA, Walker C & MacKendrick T. Pregnancy outcome among women exposed to pesticides through work or residence in an agricultural area. *Journal of Occupational Medicine* 1993 **35** 943–949. (<https://doi.org/10.1097/00043764-199309000-00019>)
- 77 Younglai EV, Foster WG, Hughes EG, Trim K & Jarrell JF. Levels of environmental contaminants in human follicular fluid, serum, and seminal plasma of couples undergoing in vitro fertilization. *Archives of Environmental Contamination and Toxicology* 2002 **43** 121–126. (<https://doi.org/10.1007/s00244-001-0048-8>)
- 78 Meeker JD, Maity A, Missmer SA, Williams PL, Mahalingaiah S, Ehrlich S, Berry KF, Altshul L, Perry MJ, Cramer DW, *et al.* Serum concentrations of polychlorinated biphenyls in relation to in vitro fertilization outcomes. *Environmental Health Perspectives* 2011 **119** 1010–1016. (<https://doi.org/10.1289/ehp.1002922>)
- 79 Milesi MM, Alarcón R, Ramos JG, Muñoz-de-Toro M, Luque EH & Varayoud J. Neonatal exposure to low doses of endosulfan induces implantation failure and disrupts uterine functional differentiation at the pre-implantation period in rats. *Molecular and Cellular Endocrinology* 2015 **401** 248–259. (<https://doi.org/10.1016/j.mce.2014.11.028>)
- 80 Guney M, Oral B, Demirin H, Take G, Giray SG, Altuntas I & Mungan T. Fallopian damage induced by organophosphate insecticide methyl parathion, and protective effect of vitamins E and C on ultrastructural changes in rats. *Toxicology and Industrial Health* 2007 **23** 429–438. (<https://doi.org/10.1177/0748233707076773>)
- 81 Paoloni-Giacobino A. Epigenetic effects of methoxychlor and vinclozolin on male gametes. In *Vitamins and Hormones*, Vol. 94, pp. 211–227. Cambridge, MA, USA: Academic Press, 2014. (<https://doi.org/10.1016/B978-0-12-800095-3.00008-0>)
- 82 Baligar PN & Kaliwal BB. Carbofuran induced block of compensatory ovarian growth in hemicastrated albino mice. *Toxicology* 2004 **204** 87–95. (<https://doi.org/10.1016/j.tox.2004.06.027>)
- 83 Guerrero-Bosagna C & Skinner MK. Environmentally induced epigenetic transgenerational inheritance of male infertility. *Current Opinion in Genetics and Development* 2014 **26** 79–88. (<https://doi.org/10.1016/j.gde.2014.06.005>)

Received in final form 8 April 2020

Accepted 7 May 2020

Accepted Manuscript published online 7 May 2020