

Table S2. Summary of previously reported composite PCC/PGL-GN with follow-up data

| Number of cases | Follow-up (time)   | Reference              |
|-----------------|--|------------------------|
| 4               | 2 patients live without disease (4-17 months);<br>1 patient died from metastasis;<br>1 patient died from hypertensive crisis during<br>resection of urothelial carcinoma | Lam et al, 1999        |
| 1               | Live without disease (11 months)   | Dundr et al, 2003      |
| 1               | Live without disease (12 months)   | Usuda et al, 2005      |
| 1               | Live without disease (8 months)  | Choi et al, 2006       |
| 1               | Live without disease (18 months)   | Hirasaki et al, 2009   |
| 1               | Live without disease   | Comstock et al, 2009   |
| 1               | Live without disease (12 months)   | Chen et al, 2009       |
| 1               | Live without disease (10 months)   | Shankar et al, 2010    |
| 1               | Live without disease   | Khan et al, 2010       |
| 1               | Live without disease (6 months)  | Menon et al, 2011      |
| 1               | Live without disease (12 months)   | Ohtsuki et al, 2012    |
| 1               | Live without disease   | Hu et al, 2013         |
| 2               | 1 patient live without disease (24 months);<br>1 patient with unknown follow-up  | Shida et al, 2015      |
| 8               | 6 patients live without disease (1-47 months);<br>2 patients died from unrelated causes  | Gupta et al, 2017      |
| 1               | Live without disease (3 months)  | Delgado et al, 2019    |
| 2               | Live without disease (19-25 months)  | Dhanasekar et al, 2021 |
| 9               | Live without disease (13-91 months)  | Our cohort             |