

Supplemental files

Supplemental table 1: Characteristics of included and excluded participants.

	Included	Excluded	P-value
n	1525	3539	
Women, n (%)	912 (59.8)	1795 (50.7)	<0.001
Age, years	57.7 ± 10.3	57.8 ± 10.6	0.861
Body mass index, kg/m ²	25.4 ± 4.1	26.6 ± 4.8	<0.001
Body mass index status, n (%)			<0.001
Normal	784 (51.4)	1396 (40.2)	
Overweight	563 (36.9)	1397 (40.2)	
Obese	178 (11.7)	681 (19.6)	
Smoking status, n (%)			<0.001
Never	692 (45.4)	1343 (38.6)	
Former	594 (39.0)	1289 (37.0)	
Current	239 (15.7)	850 (24.4)	
Physical activity, n (%)			<0.001
First	553 (36.3)	841 (31.7)	
Second	531 (34.8)	863 (32.5)	
Third	441 (28.9)	952 (35.8)	
Sedentary status, n (%)	843 (55.3)	1562 (58.8)	0.026
Hypertension, n (%)	565 (37.1)	1531 (43.4)	<0.001
Dyslipidemia, n (%)	317 (20.8)	965 (27.3)	<0.001
History of cardiovascular disease, n (%)	57 (3.7)	208 (5.9)	0.002

Results are expressed as average standard deviation for continuous variables or as number of participants (percentage) for categorical variables. Comparison between included and excluded participants performed using student's t-test for continuous variables and chi-square for categorical variables.