

**Supplemental Table 1. Characteristics and findings according to presence of distant metastasis at diagnosis (n=174).**

	Distant metastasis (-), n=116		Distant metastasis (+), n=58		P value
	Available data	Value *	Available data	Value *	
Age at diagnosis (years)	116	51.5 (40.0-63.0)	58	50.0 (40.8-67.0)	0.691
BMI (kg/m <sup>2</sup> )	88	22.8 (21.0-25.4)	43	23.3 (20.8-24.8)	0.893
Male	116	47 (40.5)	58	35 (60.3)	0.014
Initial symptom (yes)	68	52 (76.5)	38	33 (86.8)	0.199
Functional tumor (yes)	83	48 (57.8)	31	21 (67.7)	0.335
Lymph node metastasis (yes)	2	2 (100)	56	18 (32.1)	0.115
Remission (yes)	102	87 (85.3)	47	4 (8.5)	< 0.001
Recurrence (yes)	87	41 (47.1)	4	1 (25.0)	0.621
CT					
Tumor size (mm)	111	76 (55-120)	55	100 (68-140)	0.007
Pre-contrast HU	48	34.2 (30.6-39.0)	20	34.4 (31.6-39.5)	0.701
Complete adrenalectomy	114	98 (86.0)	30	27 (90.0)	0.764
Pathology					
Weiss score	50	4 (2.8-6.0)	10	5.5 (3.8-7.3)	0.042
Tumor size (mm)	106	94 (60-130)	26	130 (82.3-163.8)	0.016
Ki67 index	26	7 (4.5-16.25)	3	10 (3-20)	0.677
Mitotic count (/50HPF)	71	8 (4-21)	21	25 (7.5-57.5)	0.001
Death (yes)	81	19 (23.5)	45	29 (64.4)	< 0.001

\* Continuous variables are presented as median (interquartile range), categorical variables as n (%).

Chi-square test, Fischer's exact test, and Mann-Whitney test were adopted for comparison.